

Effect of Titrated Exposure to Non-Traumatic Noise on Unvoiced Speech Recognition in Human Listeners with Normal Audiological Profiles

Trends in Hearing
Volume 26: 1–15
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: [10.1177/23312165211117081](https://doi.org/10.1177/23312165211117081)
journals.sagepub.com/home/tia



Mengchao Zhang¹ , Richard M. Stern², Deborah Moncrieff³, Catherine Palmer⁴ and Christopher A. Brown⁴

Abstract

Non-traumatic noise exposure has been shown in animal models to impact the processing of envelope cues. However, evidence in human studies has been conflicting, possibly because the measures have not been specifically parameterized based on listeners' exposure profiles. The current study examined young dental-school students, whose exposure to high-frequency non-traumatic dental-drill noise during their course of study is systematic and precisely quantifiable. Twenty-five dental students and twenty-seven non-dental participants were recruited. The listeners were asked to recognize unvoiced sentences that were processed to contain only envelope cues useful for recognition and have been filtered to frequency regions inside or outside the dental noise spectrum. The sentences were presented either in quiet or in one of the noise maskers, including a steady-state noise, a 16-Hz or 32-Hz temporally modulated noise, or a spectrally modulated noise. The dental students showed no difference from the control group in demographic information, audiological screening outcomes, extended high-frequency thresholds, or unvoiced speech in quiet, but consistently performed more poorly for unvoiced speech recognition in modulated noise. The group difference in noise depended on the filtering conditions. The dental group's degraded performances were observed in temporally modulated noise for high-pass filtered condition only and in spectrally modulated noise for low-pass filtered condition only. The current findings provide the most direct evidence to date of a link between non-traumatic noise exposure and supra-threshold envelope processing issues in human listeners despite the normal audiological profiles.

Keywords

noise exposure, spectrotemporal envelope processing, speech in noise, cochlear synaptopathy

Received 16 December 2021; Revised received 5 July 2022; accepted 15 July 2022

Introduction

For many decades, noise exposure has been assumed to be safe or ‘non-traumatic’ if the noise does not lead to permanent threshold shifts (Saunders et al., 1985; Eggermont, 2017). However, accumulating evidence over the past ten years indicates that non-traumatic noise exposure could lead to auditory pathophysiological changes that are undetected by routine audiological exams (Kujawa & Liberman, 2009; Lin et al., 2011; Furman et al., 2013; Valero et al., 2017; Munguia et al., 2013; Sheppard et al., 2017; Pienkowski & Eggermont, 2009; Pienkowski & Eggermont, 2010a, 2010b; Pienkowski et al., 2011; Zhou & Merzenich, 2012; Fernandez et al., 2020). These physiological changes could occur at the auditory-nerve level,

¹Audiology Department, School of Life and Health Sciences, Aston University, Birmingham, B4 7ET, UK

²Department of Electrical and Computer Engineering, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

³School of Communication Sciences and Disorders, University of Memphis, Memphis, Tennessee 38152, USA

⁴Department of Communication Science and Disorders, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA

Corresponding Author:

Mengchao Zhang, Audiology Department, School of Life and Health Sciences, Aston University, Birmingham, B4 7ET, UK.
Email: zhangm22@aston.ac.uk



such as damage to the synapses of a selective group of auditory nerve fibers responsible for encoding high-intensity sound (termed ‘cochlear synaptopathy’; Kujawa & Liberman, 2009; Lin et al., 2011; Furman et al., 2013), and/or occur along the central auditory pathway, such as increase spontaneous activity (Pienkowski & Eggermont, 2009; Munguia et al., 2013; Pienkowski et al., 2013), reduced number of inhibitory neurons (Zhou & Merzenich, 2012; Munguia et al., 2013; Kamal et al., 2013; Lau et al., 2015), broadened frequency tuning curves (Zhou et al., 2011; Zhou & Merzenich, 2012; Kamal et al., 2013), reduced adaption of firing rates to sound level statistics (Bakay et al., 2018), and disrupted cortical tonotopic representation (Pienkowski & Eggermont, 2009, 2010a, 2010b; Pienkowski et al., 2011, 2013).

The current study examines one of the plausible perceptual effects of non-traumatic noise exposure, degraded supra-threshold envelope processing, which is thought to be a consequence of cochlear synaptopathy (Bharadwaj et al., 2014; Shaheen et al., 2015) though the contribution from the central auditory system is not ruled out. There has been a great deal of debates on the relationship between noise exposure and envelope processing as well as other types of auditory processing in human listeners, because the evidence has been rather inconsistent (Kumar et al., 2012; Stone et al., 2008; Stone & Moore, 2014; Paul et al., 2017, 2018; Prendergast et al., 2017a, 2017b; Yeend et al., 2017; Füllgrabe et al., 2020). It may be that human auditory perception is less susceptible to the noise exposure that does not cause permanent threshold shifts, or that researchers have overlooked consequential details in the measures or the participants. For instance, conventional tasks assessing temporal envelope processing like amplitude modulation (AM) detection or discrimination (e.g. Kumar et al., 2012; Stone et al., 2008; Paul et al., 2017) may not be always suitable for the purpose of this research, as the listeners may employ off-frequency cues for high-intensity presentation levels. Additionally, it is possible that behavioral differences will not be observed between those who have been excessively exposed and those who have not, if the differences of their exposure dosages have not reached some critical value.

The purpose of the current study is to examine the relationship between noise exposure and envelope processing abilities in humans using a different paradigm. To facilitate the distinctiveness of noise exposure profiles between the groups, dental school students were chosen as the experimental group and their non-dental-school peers as the control group. Most of the dental-school students are young adults who are less likely to have age-related hearing loss. Dental students receive regimented, well-defined exposure to dental drill noise with acoustic energy above 4 kHz (Fernandes et al., 2006; Choosong et al., 2011) throughout their program of study. As people in the general population typically get exposed to environmental noise that is often

low-pass filtered below 2 kHz (Can et al., 2010; Bořil et al., 2012; Ramo et al., 2013; Albert & Decato, 2017), the noise exposure profiles of a group of non-dental young adults should show distinct difference from those of the dental young adults. Furthermore, the levels of the noise produced by modern-day dental devices do not exceed 85 dB SPL (Chen et al., 2013; Ai et al., 2017; da Cunha et al., 2017; Ahmed & Ali, 2017; Fernandes et al., 2006; Yousuf et al., 2014; Goswami et al., 2017) so the 8-h daily time-weight average (TWA) level of dental students range between 70 and 80 dB SPL (Choosong et al., 2011; Burk & Neitzel, 2016) below legislative standards (85 dBA for 8 h, NIOSH, 1998). Taken together, these factors make dental school students a viable population to study non-traumatic noise exposure.

The current study uses recognition of unvoiced speech as a measure to assess envelope processing skills. Speech can be decomposed into temporal envelopes and temporal fine structures (TFS) (Rosen, 1992; Moore, 2019). When the TFS is replaced by random noise, speech becomes unvoiced (Kawahara & Irino, 2005; Kawahara et al., 2009) and speech intelligibility depends solely on the spectro-temporal modulations (i.e., envelopes) of the original speech. Like natural speech, intelligibility of unvoiced speech avoids potential interference by off-frequency listening because each frequency region contributes to speech differently. Despite the removal of TFS, unvoiced speech can still be highly redundant both acoustically and linguistically (Shannon et al., 1995; Loizou et al., 1999; Tillery et al., 2012; Brown, 2014, 2018). Therefore, acoustic redundancy of the unvoiced speech was controlled by filtering so the stimulus spectra fell inside or outside the spectrum of dental noise. Linguistic redundancy of the speech was constrained using speech materials with low contextual cues. Lastly, given that behavioral studies have primarily focused on temporal aspects of envelope processing (e.g. Bharadwaj et al., 2014), it is of interest whether noise exposure impacts spectral aspects of envelope processing as well. By adding temporally or spectrally modulated noise, the study examined the exposure impact on temporal or spectral envelope processing, respectively.

Routine audiological screening results and the recognition of unvoiced speech in quiet were also assessed to examine whether (1) the experimental and control groups will perform differently on unvoiced speech perception in quiet, and (2) whether non-traumatic noise exposure is related to supra-threshold envelope processing in the absence of audiological relevant peripheral changes. In theory, because non-traumatic noise exposure is thought to impact supra-threshold auditory perception, it should not impact speech perception in quiet and audiological screening outcomes. We tested these hypotheses by examining the two groups on measures of pure tone hearing sensitivity, acoustic reflexes, otoacoustic emissions, and unvoiced speech recognition in quiet.

Methods

Participant Screenings

Participants were recruited through recruitment ads, emails, and the Pitt + Me participant recruitment service sponsored by the University of Pittsburgh. All research protocols were approved by the Human Research Protection Office at the University of Pittsburgh and all participants provided written consent of participation. Before attending the formal test, participants passed demographic and audiological screenings, and a task familiarization session.

Demographic Screening. Participants filled out a questionnaire regarding their demographic information and noise exposure history. Specifically, eligible participants met the following criteria: 1) between 22 and 30 years of age; 2) speaking English as the first and the only language; 3) no known hearing issues in the past or at the time of screening, including otologic disorders, ear infection, otitis media, or hearing loss; 4) no neurological, neurophysiological or neuropsychological disorders, and no brain trauma; 5) perception of tinnitus allowed for the experimental group only if it occurred since starting dental school; 6) no history of frequent exposure to impulsive noise; 7) occupational noise exposure cannot exceed NIOSH standard (85 dBA for 8 h, NIOSH, 1998).

Participants' lifetime noise exposure dose was determined by a questionnaire that surveys the frequencies and the sound levels of noisy activities. The Exposed Noise and Hearing Disorders of Conscripts (ENHDC) questionnaire was chosen as it includes the wide range of noisy activities and allows the calculation of lifetime noise exposure dose in unit of dB SPL, informing whether the participant's past exposure has been non-traumatic. The ENHDC was first developed by Jokitalppo et al. (2006) and was revised to fit the goals of this study to include more noisy activities and collect the exposure schedule over a finer time scale, including the past 1-year, 3-year, as well as the rest of the participants' life. The activities reported are listed in Table 1. For Parts 1 to 3, all participants reported on: (1) the estimated loudness from 1 to 5 which was based on the criteria described in Jokitalppo et al., (2006), (2) the number of hours per week participating in the activity, (3) the number of weeks of participation in the past 1 year (52 weeks), (4) the number of weeks of participation from 3 years ago up to 1 year ago (104 weeks), (5) the number of years of participation and number of weeks of participation per year from birth up to 3 years ago, and (6) percentage of time using hearing protection devices. Additionally, the dental students completed Part 4 on the use of each dental drill device for each of the four school years, including (1) the number of hours per week using the device, (2) the number of weeks using the device, and (3) percentage of time using hearing protection devices. The sound levels of the dental devices were measured by the first author at the dental school. The

Table 1. Noisy Activities in the Noise Exposure Survey.

Sections and activities in the noise exposure survey

Part I – Leisure time noise

- Watching TV
- Playing video/computer games
- Working out to music
- (when not working out) Listening to music, radio programs, etc. using personal headsets or earphones
- (when not working out) Listening to music, radio programs, etc. from audio speakers in a car or at home
- Watching movies in a theatre
- Going to bars or pubs
- Attending concerts and festivals with acoustic system (e.g. classical music)
- Attending concerts, events and festivals with amplified system (e.g. rock, pop, rally)
- Attending motor sports or ride/operate motorized vehicles such as motorcycles, jet skis, speed boats, snowmobiles, or four-wheelers
- Using tools indoors during unpaid time
- Using tools outdoors during unpaid time
- Attending or participating in indoor commercial/high-school sports events (e.g. ice-hockey, basketball)
- Attending or participating in outdoor commercial/high-school sports events (e.g. football, baseball)
- Attend car/truck races
- Ride in or pilot small aircraft/private airplanes

Part 2 Leisure time noise

- Playing in a band or orchestra or singing in choir
- Practicing a musical instrument or vocal

Part 3 Occupational noise (non-dentistry noise)

- Any work involving power tools, chainsaws, or other shop tools
- Any work using drive heavy equipment or loud machinery (such as tractors, trucks, or farming or lawn equipment like mowers/leaf blowers)

Part 4 Dental noise (completed only by the dental students)

- Student handpieces
 - High-speed turbine, 82.3 dB (A)
 - Contra-angle handpiece, 70.3 dB (A)
 - Straight handpiece, 69.3 dB (A)
- Clinic handpieces
 - Ultrasonic scaler, 73.6 dB (A)
 - High-speed turbine, 80.7 dB (A)
 - Contra-angle handpiece, 62.3 dB (A)
 - Straight handpiece, 62 dB (A)
- Lab-and-clinic equipment
 - Polishing equipment, 82 dB (A)
 - Vibrating equipment, 88 dB (A)
 - Lathe equipment 3000, 93 dB (A)
 - Stone trimmer, 82.3 dB (A)
 - Low-volume suction pump, 68.3 dB (A)
 - High-volume suction pump, 69.8 dB (A)
 - Air-water syringe 60.7 dB (A)
 - Sandblaster, 90 dB (A)

dental devices were operated by a 3rd-year dental student and the levels were measured 6 inches away from the operating device using a Larson Davis 824 Sound Level Meter

(Larson Davis Inc, Depew, NY). Three measurements were performed for each device and the average levels were used in Part 4 of the noise exposure survey. The outcomes of the recording are provided in Table 1.

With the schedules and the levels available for each activity throughout a person's life, lifetime equivalent sound exposure level (L_{eq}) can be calculated. Expressing L_{eq} in dB SPL allows us to make straightforward decisions about whether a participant's noise exposure has been non-traumatic.

EQ. 1 was used to calculate the exposure dose of noisy activities (NIOSH, 1998; Neitzel et al., 1999):

$$D_i = \frac{C_i}{(8760 * N_i) / 2^{(L_i - 79)/3}} \times 100 \quad (EQ.1)$$

Where N is the number of years out of which the noise exposure needs to be computed, C is the number of hours participating in that activity during the time specified by number of years (i.e., N), L is the average sound pressure level of that activity, and i represents the ordinal number of each noisy activity. The number of hours participating in regular, less noisy activities was calculated by subtracting the total hours of noisy activity from the total hours of a participant's lifetime. These activities were assumed to occur at 64 dB SPL on average (Johnson et al., 2017). Likewise, the dose for regular activity was computed using EQ. 1. For dental-school participants, the total hours and the exposure dose of regular activity were re-calculated by subtracting the total hours of dental and non-dental noisy activities from total hours of the individual's lifetime.

The doses of all activities were then summed and used to compute L_{eq} using EQ. 2:

$$L_{eq} = \left[10 \times \log_{10} \left(\frac{\sum D_i}{100} \right) \right] + 79 \quad (EQ.2)$$

Audiological Screening. Participants received standard audiological assessments (Table 2) together with distortion product otoacoustic emissions (DPOAEs) and pure-tone audiometry at extended-high frequencies (EHFs, > 8 kHz) which may be more sensitive to noise exposure (Liberman et al., 2016). The audiologic screening included otoscopic exam, tympanometry, acoustic reflex, DPOAE, and pure-tone audiogram. The otoscopic exam was conducted at both ears using a handheld Welch Allyn otoscopy. Tympanogram was tested at both ears with a 226-Hz tone presented through a testing probe from GSI Tymistar Middle Ear Analyzer (Grason-Stadler Inc., Milford, NH).

Acoustic reflex was also conducted using GSI Tymistar Middle Ear Analyzer. Ipsilateral and contralateral acoustic reflexes were measured at each ear with a probe tone of 0.5, 1, 2, or 4 kHz presented at 95 dB SPL. The responses (in ml) of ipsilateral stimulation at the left ear were used to determine eligibility (Table 2) because the stimuli in the unvoiced speech tests were only presented at the left ear.

DPOAE was measured through Intelligent Hearing System (IHS, Miami, FL). The frequency range of the DPOAE spanned from 0.5 to 20 kHz with 3 frequency points per octave. The F2/F1 ratio was 1.22. The presentation levels for L1 and L2 were 65 and 55 dB SPL, respectively. Eligibility was determined based on the SNRs from 1 to 8 kHz (Table 2), and the SNRs from 8 to 16 kHz were used to analyze the effect of dental noise exposure on high-frequency hearing.

For the audiometric screening, pure tones ranged from 0.25 to 8 kHz (Table 2). A Madsen Astera 2 Audiometer (GN Otometrics, Denmark) controlled through the Otosuite™ software was used to present the tones over a pair of ER-3 insert earphones to the participants. The participants sat inside a soundproof booth and pressed a handheld

Table 2. Exams, Devices, and Passing Criteria of the Audiological Screening.

Exams	Devices	Passing criteria
Otoscopic exam, both ears	Handheld Welch Allyn otoscopy	No occlusion, intact ear drum
Tympanometry to 226-Hz tone, both ears	GSI Tymistar Middle Ear Analyzer	- Compliance between 0.3 ml to 1.8 ml - Middle ear pressure between -150 daPa to +150 daPa - Ear canal volume between 0.6 cc to 2.0 cc
Ipsilateral and contralateral acoustic reflexes to probe tones of 0.5, 1, 2, and 4 kHz presented at 95 dB SPL, both ears	GSI Tymistar Middle Ear Analyzer	Reflex response ≥ 0.02 ml for ipsilateral stimulation at left ear
DPOAE at f2 = 552, 698, 879, 1104, 1392, 1753, 2207, 2783, 3506, 4419, 5566, 7012, 8838, 11133, 14028, 17671 Hz, f2/f1 = 1.22, L1 = 65 dB SPL and L2 = 55 dB SPL, both ears	IHS	SNR ≥ 6 dB for 80% of the test points between 1 and 8 kHz
Pure-tone audiogram at 0.25, 0.5, 1, 2, 3, 4, 6, 8, 12.5, 14, and 16 kHz, both ears	Madsen Astera 2 with Otosuite™	Thresholds ≤ 20 dB HL from 0.25 to 8 kHz (ANSI, 2004)

bottom to indicate response to the tone. Absolute thresholds were not searched for the screening frequencies. Participants who were able to hear at or below 20 dB HL at each frequency were considered eligible. Additionally, absolute thresholds were searched and recorded for the EHF tones at 12.5, 14, and 16 kHz at each ear with Sennheiser HD 800 headphones, and the thresholds at the EHFs were further analyzed. At each frequency, a tone was presented at 25 or 30 dB HL, decreased by 10 dB if it was heard, or increase by 5 dB if no response was given. The absolute threshold was defined as the lowest level where participants gave 2 out of 3 correct responses.

Participant Information

Due to lack of previous reports using unvoiced speech recognition, a pilot study (EXP group, n = 9; CTL group, n = 7) was conducted, showing that effect size ranged from medium-large (Cohen's $f = 0.37$) to large (Cohen's $f = 0.61$) (Cohen, 1988). The total sample size for analysis of variance (ANOVA) for main effects and interactions was then estimated using G*Power 3.1 ($\alpha = 0.05$, $1 - \beta = 0.8$) based on the pilot data, yielding a total sample size between 29 to 74 participants. Fifty-two of the originally recruited eighty-six participants (EXP group, n = 25; CTL group, n = 27) passed the demographic and audiological screenings and completed the experimental speech recognition tasks. Table 3 shows means and standard deviations of age, the number of years of musical training, and lifetime non-dental noise exposure L_{eq} , lifetime dental noise exposure L_{eq} , and lifetime all noise exposure combined L_{eq} for both groups. A one-way ANOVA was conducted for each outcome variable. The lifetime L_{eq} with dental and non-dental noise combined was significantly higher for the EXP group than for the CTL group by about 3.6 dB, $F(1, 50) = 8.240$, $p = 0.006$. When considering the lifetime non-dental noise exposure L_{eq} , there was no significant

difference between the two groups, $p > 0.05$. There was also no significant difference between the two groups in age or in the years of musical training, $p > 0.05$.

For the EXP group, the lifetime dental noise exposure L_{eq} systematically and significantly increased with the number of years at dental school, where the L_{eq} of the 2nd, the 3rd, and the 4th year students were 71.2 dB SPL (SD = 1.4), 74.4 dB SPL (SD = 3.2), 77.1 dB SPL (SD = 2.6), respectively. Only 3 of 25 participants in the EXP reported the use of earplugs when they were using the student handpieces. Three different participants in the EXP group reported experience with ringing in the ear, with the tinnitus occurring either intermittently, randomly or at night.

Stimuli

Target stimuli were IEEE sentences (IEEE, 1969) that were spoken by an adult female in standard English (sampling frequency 44.1 kHz, bandwidth 0.08 to 12 kHz). An unvoiced version of each token was produced by the TANDEM-STRAIGHT vocoder (Kawahara & Irino, 2005; Kawahara et al., 2009). The vocoder extracts the envelopes of the natural utterance and excites the envelopes with random noise, producing an unvoiced token with high spectral resolution (Kawahara et al., 2009). The unvoiced stimuli were low-pass (LPF) or high-pass filtered (HPF) using a 40th-order Butterworth infinite impulse response (IIR) filter. Based on pilot data, the cut-off frequencies of the narrowest bandwidth to achieve 90% intelligibility used cutoff frequencies of 2.3 kHz for the LPF condition and 1.7 kHz for the HPF condition.

Four different maskers were used, all of which were derived from Gaussian white noise that was spectrally shaped to match the long-term average spectrum of the IEEE sentences (Figure 1, left panel). The maskers were either an unmodulated noise (UN), one of two temporally modulated noises (TMN) (Figure 1, right panel), or a

Table 3. Demographics of the EXP and the CTL Groups.

	EXP group	CTL group	F statistics
N	25 (female, n = 15)	27 (female, n = 25)	
Population	2nd to 4th year dental students (3rd to 4th year n = 20)	Non-dental graduate students or professionals with at least bachelor's degrees	
Age (years)	25.3 ± 1.7	24.6 ± 2.1	1.847
Lifetime non-dental noise exposure L_{eq} (dB SPL)	75.3 ± 4.3	75.1 ± 5.6	0.028
Lifetime dental noise exposure L_{eq} (dB SPL)	74.6 ± 3.4	0	13164.561***
Lifetime all noise exposure combined L_{eq} (dB SPL)	78.7 ± 2.7	75.1 ± 5.6	8.240**
Music training (years)	2.7 ± 3.8	3.4 ± 4.7	0.292

Note. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.




Figure 1. Left panel: spectra of the full-band unmodulated noise (solid), LPF SMN (dashed), HPF SMN (dotted). Right panel: time-domain waveforms of the full-band unmodulated noise (black), 16-Hz LPF TMN (top, grey), 16-Hz HPF TMN (bottom, grey).

Table 4. SMN Bands with Energy.

Overall filtering	Bands with energy (Hz)
Low pass	80–198
	360–585
	894–1322
	1913–2729
High pass	1913–2729
	3856–5413
	7562–10530

spectrally modulated noise (SMN). The parameters of the temporal and spectral gaps were determined by a pilot study to produce significant masking release. The TMNs were produced by sinusoidally amplitude-modulating the UN at 16 Hz or 32 Hz with a modulation depth of 1. The SMN contained spectral gaps that were 3 equivalent rectangular bands (ERB) wide and interleaved with passbands which were also 3 ERBs wide. The SMNs were processed by passing the UN through a bank of 40th-order Butterworth IIR band-pass filters. The frequencies of the unfiltered energy in the SMNs are listed in Table 4. All maskers were LPF or HPF in the same manner as the targets. The filtered maskers were then equated to the filtered targets in root-mean-square (RMS) levels.

Procedure

All psychophysical procedures were conducted in a soundproof booth. Stimulus presentation was controlled through MATLAB scripts on a MacBook Pro and presented monaurally to the left

ear through a pair of AKG K240 MKII supra-aural headphones. The exposure is assumed to have equal effect on both sides, so testing either ear should not make a difference. However, binaural presentation activates the contralateral efferent suppression on the auditory nerves (Lisowska et al., 2008). Hence, monaural presentation was used to exclude the contribution from contralateral efferent system. Before formal testing, participants performed a familiarization task in which they were instructed to repeat forty filtered unvoiced sentences (20 for each filtering condition) in quiet with feedback. Those who scored less than 90% were given an additional 10 sentences in that filtering condition. Participants who could not score 80% were excluded from the study. All participants in the current study have scored 80% or more.

The first task was to recognize unvoiced speech in quiet. The lowest sound pressure level to achieve 50% correct responses (i.e., absolute speech recognition threshold [ASRT]) was measured through a one-down-one-up adaptive procedure (Levitt, 1971) which tracks the point for 50% correct responses on the psychometric function. The initial presentation level was 0 dB SPL where participants cannot perceive the target sentence. The level was then elevated if the participant gave an incorrect response or was reduced if the participant gave a correct response. The first sentence was repeated until the participant gave a correct response, and the rest of the sentences were presented only once. The step size was 4 dB initially and 2 dB after two reversals. A correct response required correctly identifying three or more key words. The omission of the ending ‘s’ was counted correct but the omission of ‘ed’ to indicate past tense or any phoneme substitution was scored as incorrect. Three IEEE lists (i.e., 30 sentences) were used for each filtering condition




Figure 2. Tested conditions for unvoiced speech recognition task.

and the order of the filtering conditions was randomized. The measurement for a condition was stopped if the participant reached 10 reversals or completed 30 sentences, whichever came first. The ASRT was the average sound pressure levels at all but the first 2 reversals. No feedback was provided.

The second task was to recognize unvoiced speech in noise. The SNR for 50% correct responses (i.e., speech recognition threshold [SRT]) was measured through a one-down-one-up adaptive procedure. The noise level was fixed at 65 dB SPL and the target sentence level was adaptively varied. The first sentence was presented at -4 dB SNR and was repeated until the participant gave a correct response. The rules of scoring, step-size changing, and condition stopping were identical to those used in the speech in quiet task. Eight conditions (2 filtering conditions \times 4 noise maskers) were tested (Figure 2) and the order of the conditions was randomized. Three IEEE lists were used for each condition and the SRT was the average SNR of all but the first 2 reversals.

All the data were analyzed in IBM SPSS® Statistics 26.0 and plotted in MATLAB.

Results

First, audiological screening outcomes and unvoiced speech recognition performance in quiet were compared between the two groups. Next, unvoiced speech recognition in noise was compared for temporally and spectrally modulated noises. Lastly, correlation and regression analyses were conducted to examine the contributions of demographic factors and audiological screening outcomes.

Audiological Screenings and Unvoiced Speech Recognition in Quiet

The average amplitudes of the middle ear acoustic reflex for probe frequencies from 0.5 to 4 kHz were compared between the two groups through one-way analysis of variance (ANOVA). The result showed that the reflex amplitude of the CTL group (0.12 ± 0.08 ml) was on average larger than




Figure 3. Performance of LPF and HPF unvoiced speech in quiet between the CTL (dark) and the EXP (light) groups. Error bars: the standard error of the mean (SEM).

that of the EXP group (0.09 ± 0.05 ml) but the difference was not statistically significant, $F(1,50) = 2.266$, $p = 0.139$.

The DPOAE amplitudes were analyzed through a 2 (group) \times 3 (frequency) mixed-model ANOVA. Low, middle, and high frequency responses of the DPOAE were the average emission amplitudes from 0.55 to 2.8 kHz, from 3.5 to 8.8 kHz, and from 11.1 to 14 kHz, respectively. There was a significant main effect of frequency after Greenhouse-Geisser correction, $F(1.748, 87.380) = 105.025$, $p < 0.001$, $\eta_p^2 = 0.677$, but no significant main effect of group or interaction between group and frequency, $p > 0.05$. The simple-effect multiple comparisons did not show a statistically significant group difference at any given frequency condition, $p > 0.05$.

The average thresholds of pure-tone audiogram at 12.5, 14, and 16 kHz (i.e., EHF) were analyzed through a one-way ANOVA with Bonferroni correction. The thresholds of the CTL group (6.6 ± 10.9 dB HL) were slightly lower than that of the EXP group (9.2 ± 10.1 dB HL) but the difference was not statistically significant, $F(1,50) = 0.792$, $p = 0.378$.

The ASRTs for unvoiced speech recognition in quiet were analyzed through a 2 (group) \times 2 (filtering) mixed-model ANOVA (Figure 3). There was no significant main effect




Figure 4. Difference between the CTL (dark grey) and the EXP (light grey) groups in SRTs of unvoiced speech in UN (0 Hz), 16-Hz and 32-Hz TMNs under LPF (left panel) and HPF (right panel) conditions. Error bars: SEM. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

of filtering, $F(1, 50) = 3.637$, $p = 0.062$, $\eta_p^2 = 0.068$, of group, $F(1, 50) = 2.757$, $p = 0.103$, $\eta_p^2 = 0.052$, or interaction between filtering and group, $F(1) = 0.360$, $p = 0.551$, $\eta_p^2 = 0.007$ (Figure 3). There was no significant difference in ASRT between the two groups for either filtering conditions, $p > 0.05$.

Unvoiced Speech Recognition in Noise

The SRTs of unvoiced speech recognition in noise were analyzed through a 2 (group) \times 2 (filtering) \times 4 (masker) mixed-model ANOVA. There were significant main effects of group, $F(1, 50) = 6.584$, $p = 0.013$, $\eta_p^2 = 0.116$, of filtering, $F(1, 50) = 20.292$, $p < 0.001$, $\eta_p^2 = 0.289$, and of masker, $F(1, 50) = 50.889$, $p < 0.001$, $\eta_p^2 = 0.504$. There were significant interactions between group and masker, $F(3, 150) = 2.741$, $p = 0.045$, $\eta_p^2 = 0.052$, and between filtering and masker after Greenhouse-Geisser correction, $F(2.545, 127.231) = 5.851$, $p = 0.002$, $\eta_p^2 = 0.105$. There was no significant interaction between group and filtering or across the three factors.

The effect of dental noise exposure on temporal envelope processing was examined by comparing the between-group performances when filtering and masker were controlled (Figure 4). All multiple comparisons used Bonferroni correction. The mean SRTs of the CTL group always appeared lower than those of the EXP group in all TMNs (LPF, 16-Hz: mean difference [MD] = 0.8 dB; LPF, 32-Hz: MD = 1.0 dB; HPF, 16-Hz: MD = 1.9 dB; HPF, 32-Hz: MD = 3.1 dB), but a statistically significant group difference was only observed for 32-Hz TMN, $F(1, 50) = 14.112$, $p < 0.001$, $\eta_p^2 = 0.220$. Performance was also compared between the two filtering conditions within each group (Figure 5). When the modulation rate of the noise varied from 0 Hz (unmodulated) to 16 and 32 Hz, the between-filtering SRT differences increased for the CTL group, but

not for the EXP group. The SRT differences for the CTL group were 2.3 dB for 16-Hz TMN, $F(1, 50) = 10.290$, $p = 0.002$, $\eta_p^2 = 0.171$, and 3.1 dB for 32-Hz TMN, $F(1, 50) = 19.351$, $p < 0.001$, $\eta_p^2 = 0.279$. The SRT differences for the EXP group were about 1 dB (16-Hz: MD = 1 dB; 32-Hz: MD = 0.8 dB), $p > 0.05$.

The effect of exposure on spectral envelope processing also was examined in the simple-effect analysis on performance in SMN (Figure 6). The mean SRTs of the CTL group appeared lower than those of the EXP group. The group difference was statistically significant only for LPF condition (MD = 1.5 dB), $F(1, 50) = 6.853$, $p = 0.012$, $\eta_p^2 = 0.121$, but not for HPF condition (MD = 0.9 dB), $p > 0.05$.

The SRTs for the two filtering conditions were also compared within the group. Both groups scored significantly lower SRTs for the HPF condition than for the LPF condition. The differences between the filtering conditions appeared larger for the EXP group than for the CTL group. The CTL group showed an SRT difference of 1.8 dB between the LPF and the HPF conditions, $F(1, 50) = 5.741$, $p = 0.020$, $\eta_p^2 = 0.103$. The EXP group an SRT difference of 2.3 dB, $F(1, 50) = 9.413$, $p = 0.003$, $\eta_p^2 = 0.158$.

Contributions of Demographic and Audiological (Screening) Factors

Based on Figures 4 and 5, the EXP group performed more poorly on average than the CTL group in temporally or spectrally modulated noises, though the difference for the temporal condition did not reach statistical significance. It is of interest whether the participants' demographic and audiological factors have contributed to the variations of the LPF unvoiced speech recognition. Therefore, LPF speech measures, including the SRTs of LPF unvoiced speech in




Figure 5. SRT differences between LPF (dark grey) and HPF (light grey) unvoiced speech in UN and TMNs within the CTL (left panel) and the EXP group (right panel). Error bars: SEM. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.




Figure 6. SRTs of unvoiced speech in SMN under LPF (left panel) and HPF (right panel) conditions. Error bars: SEM. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

16-Hz TMN, in 32-Hz TMN, and in SMN, were analyzed using hierarchical multiple linear regression (HMLR). The basic demographic factors included age and years of musical training. The exposure-related demographic factors included the lifetime dental noise exposure L_{eq} (in dB SPL) and the lifetime non-dental noise exposure L_{eq} (in dB SPL). The variable for middle ear function was the average acoustic reflex amplitude (in ml) of low and middle frequencies (i.e. 0.5 to 2 kHz), which covered the frequency of the LPF unvoiced speech. The variables for inner ear outer hair cell (OHC) function were the average SNRs of DPOAE at low stimulus frequencies (0.552 to 2.783 kHz) and at high stimulus frequencies (3.506 to 8.838 kHz). The sequence of the factors adding into the HMLR analysis was 1) the basic demographic factors, 2) the exposure-related factors, 3) the

variable of middle-ear function, and 4) the variable of inner ear OHC function. The EHF measures, such as EHF DPOAE and PTAs, were not correlated to the LPF speech measures. And as these measures examined the frequencies distant from the LPF speech frequencies (i.e. < 2.3 kHz), they were excluded from the regression analysis.

For LPF unvoiced speech in 16-Hz or 32-Hz TMNs, there were no significant correlations between the SRTs and any of the predictors, indicating no linear relationships between any of the demographic or audiologic factors and the SRTs of the LPF unvoiced speech in TMNs. Hence, linear regression was not conducted. For LPF unvoiced speech in SMN, the SRT was significantly correlated with lifetime dental noise exposure L_{eq} , $r = 0.337$, $p = 0.015$. Tests of normality, homoscedasticity, and multicollinearity were not violated. The HMLR

Table 5. HMLR Model for the SRT of LPF Unvoiced Speech in SMN.

	Model 1			Model 2			Model 3			Model 4		
	B	SE	β	B	SE	β	B	SE	β	B	SE	β
Age	0.054	0.162	0.047	-0.088	0.158	-0.078	-0.088	0.16	-0.077	-0.092	0.172	-0.081
Years of music training	0.060	0.074	0.116	0.061	0.07	0.117	0.06	0.074	0.116	0.079	0.079	0.152
Dental L _{eq}				0.022	0.008	0.368**	0.022	0.008	0.369*	0.021	0.008	0.366*
Non-dental L _{eq}				-0.102	0.06	-0.23	-0.102	0.063	-0.231	-0.097	0.065	-0.221
Acoustic reflex amplitude							0.19	4.812	0.006	-0.374	5.166	-0.011
DPOAE amplitude (low frequency)										0.007	0.076	0.014
DPOAE amplitude (high frequency)										-0.068	0.078	-0.124
R ²		0.017			0.185			0.185			0.199	
ΔR ²		0.017			0.168			< 0.001			0.014	

Note. *, p < 0.05; **, p < 0.01.

model was built, showing that lifetime dental noise exposure L_{eq} was the only significant predictor, while age, years of musical training, non-dental noise exposure, acoustic reflex amplitudes, or DPOAE amplitudes failed to account for the variance of the performance. The final model explained 19.9% of the variance in performance (Table 5).

Discussion

The relationship between non-traumatic noise exposure and supra-threshold auditory envelope processing in human listeners was examined here. This study utilized unvoiced speech recognition that relies solely on temporal and spectral envelopes among dental-school students with quantifiable exposure to non-traumatic noise during professional training. A between-groups design was implemented to compare dental-school students to a cohort of peers not enrolled in dental school. The two groups showed no statistically significant differences in general demographic or audiological screening outcomes. Also, no difference was observed between-group on unvoiced speech recognition when the speech was presented in quiet or in unmodulated noise. When the noise was modulated, and listeners were required to exploit the temporal or spectral gaps with favorable SNRs to improve recognition, the group with dental noise exposure performed more poorly than their unexposed peers, and the poor use of spectral or temporal cues appeared dependent on stimulus frequency. Non-traumatic exposure to high-frequency dental noise was found to be associated with poor temporal envelope processing at higher frequencies but poor spectral envelope processing at lower frequencies. Given that the sound levels from the dental equipment typically do not exceed 80 dB SPL and students do not practice for more than 8 h per day, these results support the hypothesis that non-traumatic noise exposure may contribute to the degradation of supra-threshold envelope processing of speech that cannot be detected by routine audiological screenings.

The study first found that the experience of non-traumatic noise exposure could be related to poor temporal envelope processing. Listeners with dental noise exposure did not show as much masking release as the control listeners when recognizing the HPF unvoiced speech in TMN. When modulation rate was increased from 16 to 32 Hz, masking release increased in smaller magnitude for the EXP than for the CTL listeners. At higher modulation rates, the temporal gaps of the noise become briefer, so the task demands on temporal resolution for exploiting the information in the gaps could grow accordingly (Gustafsson & Arlinger, 1994; Dubno et al., 2003) and accentuate the poor temporal resolution of the compromised auditory system.

One of the plausible explanations for the links between noise exposure and poor supra-threshold temporal envelope processing is cochlear synaptopathy (Bharadwaj et al., 2014; Shaheen et al., 2015). Cochlear synaptopathy is a pathological change in the synapses of a selective group of auditory nerve fibers that can be induced by exposing the individual to noise that does not induce permanent threshold shift (Kujawa & Liberman, 2009; Lin et al., 2011; Furman et al., 2013). These auditory fibers normally encode the sound envelopes or intensity changes at high sound levels, varying the firing rates according to the input level changes while the firing rates of other fibers have saturated (Liberman, 1978). If these auditory fibers are damaged, one of the consequences is thought to be the degraded encoding of sound envelopes at supra-threshold levels. The current finding may also be explained by the reduced dynamic range adaptation in the neurons of the inferior colliculus (Bakay et al., 2018). Dynamic range adaptation refers to the ability of a neuron to shift its rate-level function toward the frequently occurring sound levels to avoid firing saturation and ensure high fidelity when encoding various sound levels (Dean et al., 2008). In Bakay et al. (2018), the amount of dynamic range adaption by the inferior colliculus

neurons reduces after the noise exposure has induced cochlear synaptopathy, suggesting that the ability of temporal envelope coding could be impacted in the inferior colliculus in addition to cochlear synaptopathy.

The relation of temporal envelope processing and noise exposure has been examined previously, but the results have been conflicting. Kumar et al. (2012) found poorer AM detection thresholds for train drivers with normal audiograms than for age-matched unexposed individuals at 60 and 200 Hz modulation rates along with poor duration pattern test and speech reception in noise, supporting the temporal hypothesis of synaptopathy. Meanwhile, Paul et al. (2017, 2018) used AM detection with stimuli presented in narrow-band noise at various noise levels to limit off-frequency listening and to engage auditory fibers of different spontaneous rates. They did not consistently show poorer performance by young adults with higher noise exposure compared to their peers with lower exposure. Yeend et al. (2017) and Füllgrabe et al. (2020) controlled off-frequency listening by presenting the modulated targets in threshold-equalizing noise (TEN), which is configured to produce equal tone-in-noise thresholds for normal hearing listeners from 0.25 to 10 kHz and is used in the TEN test to diagnose cochlear dead regions (Moore et al., 2000), and they found no relationship between noise exposure and temporal processing. Stone et al. (2008) and Stone and Moore (2014) found worse AM discrimination for a noise-exposed group but only when the signals were presented near-threshold and not supra-threshold.

The lack of consistent previous evidence to associate temporal envelope processing and noise exposure suggest that conventional psychophysical measures of temporal envelope processing may not be sensitive enough or need further parameterization. This study has shown that unvoiced speech recognition in TMN can be an alternative and potentially more desirable measure to assess temporal envelope processing after non-traumatic noise exposure. Speech-based tasks provide the advantage of controlling off-frequency listening because speech intelligibility relies on the combined contributions of various frequency regions. The use of unvoiced speech can elicit high intelligibility in quiet¹ despite the removal of pitch and harmonic information, which places greater emphasis on envelope cues. Then, bandpass filtering can constrain the examination to spectral regions of interest, such as those thought to be affected by noise exposure. The addition of TMN manipulates the test toward assessing temporal resolution and the efficiency of extracting information from the temporal gaps of modulated noise can be assessed using properly selected modulation rates, as has been shown in this study.

That said, when spectrally modulated noise was added, unvoiced speech recognition was weighed toward using spectral envelope cues. The current study also discovered that the experience of non-traumatic noise exposure could be related to spectral envelope processing. Listeners with dental noise exposure performed poorly when recognizing

the LPF unvoiced speech in 3-ERB gapped SMN. The finding is not surprising. Without normal high-intensity auditory fibers, the rest of the fibers cannot fully represent the sound spectra at high intensities without firing-rate saturation, from which the reconstructed spectra may appear smoothed out with impoverished spectral details (Sachs & Young, 1979; May et al., 1996; Reiss et al., 2011). Furthermore, studies on the central auditory system have also reported negative impacts of non-traumatic noise exposure on neuronal spectral encoding, such as disrupted cortical tonotopic representation (Pienkowski & Eggermont, 2009, 2010a, 2010b; Pienkowski et al., 2011, 2013) and broadened neuronal frequency tuning curves (Zhou et al., 2011; Zhou & Merzenich, 2012; Kamal et al., 2013). Most of these studies have adopted moderate level of noise exposure (65 to 80 dB SPL) with prolonged exposure schedules (4 to 12 weeks), providing compelling evidence on the impact of prolonged non-traumatic noise exposure, such as dental noise, on spectral envelope processing.

It was interesting, however, to observe the performance difference between the two groups shifted from LPF to HPF speech when noise modulation switched from spectral into temporal domains. The exact reason behind the finding is yet unclear. One plausible explanation is that there is a trade-off between spectral and temporal resolution along the frequency axis and the degrees of spectral and temporal resolution vary from low to high frequencies. As the basilar membrane response is often modeled as a bank of Gammatone filters (Patterson et al., 1992; Lopez-Poveda & Meddis, 2001) and the impulse responses of the filters decreases in duration with increasing center frequencies, the temporal resolution of the auditory system may improve with increasing best frequencies while spectral resolution acts the opposite way. Another explanation is that the spetro-temporal trade-off along the frequency axis does not occur in the auditory system but in the importance weighting of speech cues. Previous studies using vocoded speech, which only relies on temporal envelope cues for speech recognition, have shown that despite temporal envelope cues appear important at all frequencies for speech in quiet, these cues are more heavily weighted at high frequencies than low frequencies for speech in noise (Apoux & Bacon, 2004; Ardoit et al., 2011; Fogerty, 2011).

It should be noted that the two groups did not appear different on the acoustic reflex amplitudes or the EHF measures, which were previously thought to be impacted by non-traumatic noise exposure (Valero et al., 2018; Liberman et al., 2016). Lack of reflex difference may be because that the acoustic reflex used in this study was a clinical screening, which had not controlled for spread of excitation at high stimulus intensities. Lack of the EHF threshold difference or DPOAE difference may be because of the different susceptibility to noise exposure in humans and animals. It will be interesting to examine whether group difference will emerge for the EHF measures on a long run, such as an

early onset of EHF hearing loss in noise exposed group (Fernandez et al., 2015).

In conclusion, exposure to non-traumatic noise over time could be related to reduced envelope processing in humans in the absence of clinically defined audiological abnormalities. The finding supported the general hypothesis of cochlear synaptopathy, though central auditory dysfunction could also play a role. The effect of noise exposure could be related to both spectral and temporal processing and impact the two aspects of envelope processing in a frequency-dependent fashion. The task of recognizing unvoiced speech in modulated noise is shown to be usable in revealing supra-threshold envelope processing issues. However, the current study is limited in several aspects. A cross-sectional study coupled with a within-subject longitudinal design would be ideal for observing the noise exposure impact over time. It should be noted that despite the numbers of female and male participants were less balanced for the CTL than for the EXP groups, the result patterns remained when only female participants were examined and there was no significant performance difference between the males and the females within the EXP group (data not shown). Like all psychophysical studies, the current measure does not identify the site of lesions or the physiological mechanisms behind non-traumatic noise exposure. Approaches like computational modeling of pathological conditions in physiologically inspired auditory models may be useful to parameterize the current measure to differentially assess various site of lesions. Electrophysiological measures, such as envelope following responses, may also benefit the mechanistic study relating to temporal envelope processing and noise exposure. Overall, if the goal is to early discover the impact of noise exposure on hearing before measurable hearing threshold change, it is worth considering the inclusion of unvoiced speech recognition in a proposed test battery.

Acknowledgements

We would like to thank Dr. Bharath Chandrasekaran (University of Pittsburgh) for his insightful comments to the early versions of the manuscript. This study was a part of the dissertation of the first author and was supported by NIDCD grant R21DC018408, and a PhD Student Research Award from the School of Health and Rehabilitation Sciences at the University of Pittsburgh.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the National Institute on Deafness and Other Communication Disorders, School of Health and Rehabilitation Sciences, University of Pittsburgh, (grant number R21DC018408, PhD Student Research Award).

ORCID iD

Mengchao Zhang  <https://orcid.org/0000-0002-7361-9969>

Note

1. Our pilot data ($n = 5$), young normal hearing listeners achieved near 100% correct performance for unfiltered unvoiced speech recognition.

Reference

- Ahmed H. O., & Ali W. J. (2017). Noise levels, noise annoyance, and hearing-related problems in a dental college. *Archives of Environmental & Occupational Health*, 72(3), 159–165. <https://doi.org/10.1080/19338244.2016.1179169>
- Ai Z., Mak C., & Wong H. (2017). Noise level and its influences on dental professionals in a dental hospital in Hong Kong. *Building Services Engineering Research and Technology*, 38(5), 522–535. <https://doi.org/10.1177/0143624417705529>
- Albert D. G., & Decato S. N. (2017). Acoustic and seismic ambient noise measurements in urban and rural areas. *Applied Acoustics*, 119, 135–143. <https://doi.org/10.1016/j.apacoust.2016.12.015>
- Apoux F., & Bacon S. P. (2004). Relative importance of temporal information in various frequency regions for consonant identification in quiet and in noise. *The Journal of the Acoustical Society of America*, 116(3), 1671–1680. <https://doi.org/10.1121/1.1781329>
- Ardoit M., Agus T., Sheft S., & Lorenzi C. (2011). Importance of temporal-envelope speech cues in different spectral regions. *The Journal of the Acoustical Society of America*, 130(2), EL115–EL121. <https://doi.org/10.1121/1.3602462>
- Bakay W. M. H., Anderson L. A., Garcia-Lazaro J. A., McAlpine D., & Schaette R. (2018). Hidden hearing loss selectively impairs neural adaptation to loud sound environments. *Nature Communications*, 9(1), 4298. <https://doi.org/10.1038/s41467-018-06777-y>
- Bharadwaj H. M., Verhulst S., Shaheen L., Liberman M. C., & Shinn-Cunningham B. G. (2014). Cochlear neuropathy and the coding of supra-threshold sound. *Frontiers in Systems Neuroscience*, 8, 26. <https://doi.org/10.3389/fnsys.2014.00026>
- Bofil H., Boyraz P., & Hansen J. H. L. (2012). Towards multimodal driver's stress detection. In J. H. L. Hansen, P. Boyraz, K. Takeda, & H. Abut (Eds.), *Digital signal processing for in-vehicle systems and safety* (pp. 3–19). Springer. https://doi.org/10.1007/978-1-4419-9607-7_1.
- Brown C. A. (2014). Binaural enhancement for bilateral cochlear implant users. *Ear & Hearing*, 35(5), 580–584. <https://doi.org/10.1097/AUD.0000000000000044>
- Brown C. A. (2018). Corrective binaural processing for bilateral cochlear implant patients. *PLOS ONE*, 13(1), e0187965. <https://doi.org/10.1371/journal.pone.0187965>
- Burk A., & Neitzel R. L. (2016). An exploratory study of noise exposures in educational and private dental clinics. *Journal of Occupational and Environmental Hygiene*, 13(10), 741–749. <https://doi.org/10.1080/15459624.2016.1177646>
- Can A., Leclercq L., Lelong J., & Botteldooren D. (2010). Traffic noise spectrum analysis: dynamic modeling vs. Experimental observations. *Applied Acoustics*, 71(8), 764–770. <https://doi.org/10.1016/j.apacoust.2010.04.002>

- Chen W.-L., Chen C.-J., Yeh C.-Y., Lin C.-T., Cheng H.-C., & Chen R.-Y. (2013). Workplace noise exposure and its consequent annoyance to dentists. *Journal of Experimental & Clinical Medicine*, 5(5), 177–180. <https://doi.org/10.1016/j.jecm.2013.08.009>
- Choosong T., Kaimook W., Tantisarasart R., Sooksamear P., & Chayaphum S., Kongkamol, C., ... Phakthongsuk, P. (2011). Noise exposure assessment in a dental school. *Safety and Health at Work*, 2(4), 348–354.
- Cohen J. (1988). *Statistical power analysis for the behavioral sciences*. Routledge Academic.
- da Cunha K. F., dos Santos R. B., & Klien C. A. (2017). Assessment of noise intensity in a dental teaching clinic. *BDJ Open*, 3(1), 1–3. <https://doi.org/10.1038/bdjopen.2017.10>
- Dean I., Robinson B. L., Harper N. S., & McAlpine D. (2008). Rapid neural adaptation to sound level statistics. *Journal of Neuroscience*, 28(25), 6430–6438. <https://doi.org/10.1523/JNEUROSCI.0470-08.2008>
- Dubno J. R., Horwitz A. R., & Ahlstrom J. B. (2003). Recovery from prior stimulation: masking of speech by interrupted noise for younger and older adults with normal hearing. *The Journal of the Acoustical Society of America*, 113(4), 2084–2094. <https://doi.org/10.1121/1.1555611>
- Eggermont J. J. (2017). Effects of long-term non-traumatic noise exposure on the adult central auditory system. Hearing problems without hearing loss. *Hearing research*, 352, 12–22.
- Fernandes J. C., Carvalho A. P. O., Gallas M., Vaz P., & Matos P. A. (2006). Noise levels in dental schools. *European Journal of Dental Education*, 10(1), 32–37.
- Fernandez K. A., Guo D., Micucci S., De Gruttola V., Liberman M. C., & Kujawa S. G. (2020). Noise-induced cochlear synaptopathy with and without sensory cell loss. *Neuroscience*, 427, 43–57. <https://doi.org/10.1016/j.neuroscience.2019.11.051>
- Fernandez K. A., Jeffers P. W. C., Lall K., Liberman M. C., & Kujawa S. G. (2015). Aging after noise exposure: acceleration of cochlear synaptopathy in “recovered” ears. *Journal of Neuroscience*, 35(19), 7509–7520. <https://doi.org/10.1523/JNEUROSCI.5138-14.2015>
- Fogerty D. (2011). Perceptual weighting of the envelope and fine structure across frequency bands for sentence intelligibility: effect of interruption at the syllabic-rate and periodic-rate of speech. *The Journal of the Acoustical Society of America*, 130(1), 489–500. <https://doi.org/10.1121/1.3592220>
- Füllgrabe C., Moody M., & Moore B. C. J. (2020). No evidence for a link between noise exposure and auditory temporal processing for young adults with normal audiograms. *The Journal of the Acoustical Society of America*, 147(6), EL465–EL470. <https://doi.org/10.1121/10.0001346>
- Furman A. C., Kujawa S. G., & Liberman M. C. (2013). Noise-induced cochlear neuropathy is selective for fibers with low spontaneous rates. *Journal of Neurophysiology*, 110(3), 577–586.
- Goswami M., Singh D., Vashist B., & Marwaha S. (2017). Noise levels and sound pollution associated with various operative procedures and equipments in a pediatric dental environment—A clinical study. *Journal of Oral Biology and Craniofacial Research*, 7(3), 182–187. <https://doi.org/10.1016/j.jobcr.2017.06.003>
- Gustafsson HÅ, & Arlinger S. D. (1994). Masking of speech by amplitude-modulated noise. *The Journal of the Acoustical Society of America*, 95(1), 518–529. <https://doi.org/10.1121/1.408346>
- IEEE Recommended Practice for Speech Quality Measurements. (1969). *IEEE Transactions on Audio and Electroacoustics*, 17(3). <https://doi.org/10.1109/TAU.1969.1162058>.
- Johnson T. A., Cooper S., Stamper G. C., & Chertoff M. (2017). Noise exposure questionnaire: A tool for quantifying annual noise exposure. *Journal of the American Academy of Audiology*, 28(01), 014–035. <https://doi.org/10.3766/jaaa.15070>
- Jokitalppo J., Toivonen M., & Bjoörk E. (2006). Estimated leisure-time noise exposure, hearing thresholds, and hearing symptoms of Finnish conscripts. *Military Medicine*, 171(2), 112–116. <https://doi.org/10.7205/MILMED.171.2.112>
- Kamal B., Holman C., & de Villers-Sidani E. (2013). Shaping the aging brain: role of auditory input patterns in the emergence of auditory cortical impairments. *Frontiers in Systems Neuroscience*, 7, 52. <https://doi.org/10.3389/fnsys.2013.00052>
- Kawahara H., & Irino T. (2005). Underlying principles of a high-quality speech manipulation system STRAIGHT and its application to speech segregation. In *Speech separation by humans and machines*. (pp. 167–180). Boston, MA: Springer. https://doi.org/10.1007/0-387-22794-6_11
- Kawahara H., Takahashi T., Morise M., & Banno H. (2009). Development of exploratory research tools based on TANDEM-STRAIGHT. *APSIPA ASC 2009 - Asia-Pacific Signal and Information Processing Association 2009 Annual Summit and Conference*.
- Kujawa S. G., & Liberman M. C. (2009). Adding insult to injury: Cochlear nerve degeneration after “temporary” noise-induced hearing loss. *Journal of Neuroscience*, 29(45), 14077–14085.
- Kumar U. A., Ameenudin S., & Sangamanatha A. V. (2012). Temporal and speech processing skills in normal hearing individuals exposed to occupational noise. *Noise & Health*, 14(58), 100. <https://doi.org/10.4103/1463-1741.97252>
- Levitt H. (1971). Transformed up-down methods in psychoacoustics. *The Journal of the Acoustical Society of America*, 49(2B), 467–477. <https://doi.org/10.1121/1.1912375>
- Liberman M. C. (1978). Auditory-nerve response from cats raised in a low-noise chamber. *The Journal of the Acoustical Society of America*, 63(2), 442–455. <https://doi.org/10.1121/1.381736>
- Liberman M. C., Epstein M. J., Cleveland S. S., Wang H., & Maison S. F. (2016). Toward a differential diagnosis of hidden hearing loss in humans. *PLOS ONE*, 11(9), e0162726. <https://doi.org/10.1371/journal.pone.0162726>
- Lin H. W., Furman A. C., Kujawa S. G., & Liberman M. C. (2011). Primary neural degeneration in the Guinea pig cochlea after reversible noise-induced threshold shift. *Journal of the Association for Research in Otolaryngology*, 12(5), 605–616.
- Lisowska G., Namysłowski G., & Misiołek M., Wojciech, Ś., Orecka, B., Czecior, E., & Dziendziel, A. (2008). Efferent suppression test-sensitivity and specific city. *Otolaryngologia Polska*, 62, 747–754.
- Loizou P. C., Dorman M., & Tu Z. (1999). On the number of channels needed to understand speech. *The Journal of the Acoustical Society of America*, 106(4), 2097–2103. <https://doi.org/10.1121/1.427954>
- Lopez-Poveda E. A., & Meddis R. (2001). A human nonlinear cochlear filterbank. *The Journal of the Acoustical Society of America*, 110(6), 3107–3118. <https://doi.org/10.1121/1.1416197>

- May B. J., Huang A., Prell L. E., & Hienz G., & D R. (1996). Vowel formant frequency discrimination in cats: comparison of auditory nerve representations and psychophysical thresholds. *Auditory Neuroscience*, 3(2), 135–162.
- Moore B. C. J., Huss M., Vickers D. A., Glasberg B. R., & Alcántara J. I. (2000). A test for the diagnosis of dead regions in the cochlea. *British Journal of Audiology*, 34(4), 205–224.
- Moore B. C. J. (2019). The roles of temporal envelope and fine structure information in auditory perception. *Acoustical Science and Technology*, 40(2), 61–83. <https://doi.org/10.1250/ast.40.61>
- Munguia R., Pienkowski M., & Eggermont J. J. (2013). Spontaneous firing rate changes in cat primary auditory cortex following long-term exposure to non-traumatic noise: tinnitus without hearing loss? *Neuroscience Letters*, 546, 46–50. <https://doi.org/10.1016/j.neulet.2013.04.048>
- National Institute for Occupational Safety and Health. (1998). Criteria for a recommended standard: Occupational noise exposure, revised criteria 1998. US Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention.
- Neitzel R., Seixas N. S., Camp J., & Yost M. (1999). An assessment of occupational noise exposures in four construction trades. *American Industrial Hygiene Association Journal*, 60(6), 807–817. <https://doi.org/10.1080/00028899908984506>
- Patterson R. D., Robinson K., Holdsworth J., McKeown D., Zhang C., & Allerhand M. (1992). *Complex Sounds and Auditory Images*. In *Auditory physiology and perception* (pp. 429–446). Elsevier. <https://doi.org/10.1016/B978-0-08-041847-6.50054-X>.
- Paul B. T., Bruce I. C., & Roberts L. E. (2018). Envelope following responses, noise exposure, and evidence of cochlear synaptopathy in humans: correction and comment. *The Journal of the Acoustical Society of America*, 143(6), EL487–EL489. <https://doi.org/10.1121/1.5043082>
- Paul B. T., Waheed S., Bruce I. C., & Roberts L. E. (2017). Subcortical amplitude modulation encoding deficits suggest evidence of cochlear synaptopathy in normal-hearing 18–19 year olds with higher lifetime noise exposure. *The Journal of the Acoustical Society of America*, 142(5), EL434–EL440. <https://doi.org/10.1121/1.5009603>
- Pienkowski M., & Eggermont J. J. (2009). Long-term, partially-reversible reorganization of frequency tuning in mature cat primary auditory cortex can be induced by passive exposure to moderate-level sounds. *Hearing Research*, 257(1–2), 24–40. <https://doi.org/10.1016/j.heares.2009.07.011>
- Pienkowski M., & Eggermont J. J. (2010a). Intermittent exposure with moderate-level sound impairs central auditory function of mature animals without concomitant hearing loss. *Hearing Research*, 261(1–2), 30–35. <https://doi.org/10.1016/j.heares.2009.12.025>
- Pienkowski M., & Eggermont J. J. (2010b). Passive exposure of adult cats to moderate-level tone pip ensembles differentially decreases AI and AII responsiveness in the exposure frequency range. *Hearing Research*, 268(1–2), 151–162. <https://doi.org/10.1016/j.heares.2010.05.016>
- Pienkowski M., Munguia R., & Eggermont J. J. (2011). Passive exposure of adult cats to bandlimited tone pip ensembles or noise leads to long-term response suppression in auditory cortex. *Hearing Research*, 277(1–2), 117–126. <https://doi.org/10.1016/j.heares.2011.02.002>
- Pienkowski M., Munguia R., & Eggermont J. J. (2013). Effects of passive, moderate-level sound exposure on the mature auditory cortex: spectral edges, spectrotemporal density, and real-world noise. *Hearing Research*, 296, 121–130. <https://doi.org/10.1016/j.heares.2012.11.006>
- Prendergast G., Guest H., Munro K. J., Kluk K., Léger A., Hall D. A., Heinz M. G., & Plack C. J. (2017a). Effects of noise exposure on young adults with normal audiograms I: electrophysiology. *Hearing Research*, 344, 68–81. <https://doi.org/10.1016/j.heares.2016.10.028>
- Prendergast G., Millman R. E., Guest H., Munro K. J., Kluk K., Dewey R. S., Hall D. A., Heinz M. G., & Plack C. J. (2017b). Effects of noise exposure on young adults with normal audiograms II: behavioral measures. *Hearing Research*, 356, 74–86. <https://doi.org/10.1016/j.heares.2017.10.007>
- Ramo J., Valimaki V., & Tikander M. (2013). Perceptual headphone equalization for mitigation of ambient noise. 2013 *IEEE International Conference on Acoustics, Speech and Signal Processing*, 724–728. <https://doi.org/10.1109/ICASSP.2013.6637743>.
- Reiss L. A. J., Ramachandran R., & May B. J. (2011). Effects of signal level and background noise on spectral representations in the auditory nerve of the domestic cat. *Journal of the Association for Research in Otolaryngology*, 12(1), 71–88. <https://doi.org/10.1007/s10162-010-0232-5>
- Rosen S. (1992). Temporal information in speech: acoustic, auditory and linguistic aspects. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 336(1278), 367–373. <https://doi.org/10.1098/rstb.1992.0070>
- Sachs M. B., & Young E. D. (1979). Encoding of steady-state vowels in the auditory nerve: representation in terms of discharge rate. *Journal of the Acoustical Society of America*, 66(2), 470–479. <https://doi.org/10.1121/1.383098>
- Saunders J. C., Dear S. P., & Schneider M. E. (1985). The anatomical consequences of acoustic injury: a review and tutorial. *The Journal of the Acoustical Society of America*, 78(3), 833–860.
- Shaheen L. A., Valero M. D., & Liberman M. C. (2015). Towards a diagnosis of cochlear neuropathy with envelope following responses. *Journal of the Association for Research in Otolaryngology*, 16(6), 727–745. <https://doi.org/10.1007/s10162-015-0539-3>
- Shannon R. V., Zeng F. G., Kamath V., Wygonski J., & Ekelid M. (1995). Speech recognition with primarily temporal cues. *Science*, 270(5234), 303–304. <https://doi.org/10.1126/science.270.5234.303>
- Sheppard A. M., Chen G.-D., Manohar S., Ding D., Hu B.-H., Sun W., Zhao J., & Salvi R. (2017). Prolonged low-level noise-induced plasticity in the peripheral and central auditory system of rats. *Neuroscience*, 359, 159–171. <https://doi.org/10.1016/j.neuroscience.2017.07.005>
- Stone M. A., & Moore B. C. J. (2014). On the near non-existence of “pure” energetic masking release for speech. *The Journal of the Acoustical Society of America*, 135(4), 1967–1977. <https://doi.org/10.1121/1.4868392>
- Stone M. A., Moore B. C. J., & Greenish H. (2008). Discrimination of envelope statistics reveals evidence of sub-clinical hearing damage in a noise-exposed population with “normal” hearing thresholds. *International Journal of Audiology*, 47(12), 737–750. <https://doi.org/10.1080/14992020802290543>
- Tillery K. H., Brown C. A., & Bacon S. P. (2012). Comparing the effects of reverberation and of noise on speech recognition in

- simulated electric-acoustic listening. *The Journal of the Acoustical Society of America*, 131(1), 416–423. <https://doi.org/10.1121/1.3664101>
- Valero M. D., Burton J. A., Hauser S. N., Hackett T. A., Ramachandran R., & Liberman M. C. (2017). Noise-induced cochlear synaptopathy in rhesus monkeys (*Macaca mulatta*). *Hearing Research*, 353, 213–223. <https://doi.org/10.1016/j.heares.2017.07.003>
- Valero M. D., Hancock K. E., Maison S. F., & Liberman M. C. (2018). Effects of cochlear synaptopathy on middle-ear muscle reflexes in unanesthetized mice. *Hearing Research*, 363, 109–118.
- Yeend I., Beach E. F., Sharma M., & Dillon H. (2017). The effects of noise exposure and musical training on suprathreshold auditory processing and speech perception in noise. *Hearing Research*, 353, 224–236. <https://doi.org/10.1016/j.heares.2017.07.006>
- Yousuf A., Ganta S., Nagaraj A., Pareek S., Atri M., Singh K., & Sidiq M. (2014). Acoustic noise levels of dental equipments and its association with fear and annoyance levels among patients attending different dental clinic setups in Jaipur, India. *Journal of Clinical and Diagnostic Research: JCDR*, 8(4), ZC29. <https://doi.org/10.7860/JCDR/2014/7678.4245>
- Zhou X., & Merzenich M. M. (2012). Environmental noise exposure degrades normal listening processes. *Nature Communications*, 3(1), 1–9. <https://doi.org/10.1038/ncomms1849>
- Zhou X., Panizzutti R., de Villers-Sidani E., Madeira C., & Merzenich M. M. (2011). Natural restoration of critical period plasticity in the juvenile and adult primary auditory Cortex. *Journal of Neuroscience*, 31(15), 5625–5634. <https://doi.org/10.1523/JNEUROSCI.6470-10.2011>