

**Title:** Social cognition and idiopathic isolated cervical dystonia

**Running head:** Social cognition in cervical dystonia

Kristína Czekóová<sup>1</sup>, Petra Zemánková<sup>1,2</sup>, Daniel J. Shaw<sup>1</sup>, Martin Bareš<sup>1,2,3</sup>

<sup>1</sup>CEITEC, Masaryk University, Brno, Czech Republic

<sup>2</sup>First Department of Neurology, Faculty of Medicine, Masaryk University and St. Anne's Teaching Hospital, Brno, Czech Republic

<sup>3</sup>Department of Neurology, Medical School, University of Minnesota, Minneapolis, USA

Correspondence: Kristína Czekóová, Kamenice 753/5, 62500 Brno, Czech Republic

+420 549496304; [kristina.czekoova@ceitec.muni.cz](mailto:kristina.czekoova@ceitec.muni.cz); ORCID 0000-0001-9198-1083

**Keywords:** cervical dystonia; social cognition; Theory of mind; empathy; cognition

#### **Disclosure of conflicts of interest**

Nothing to report.

#### **Funding**

This work was funded by the Ministry of Education, Youth and Sports of the Czech Republic under the project CEITEC 2020 (LQ1601).

## **Abstract**

### *Background*

For a long time cervical dystonia (CD) has been characterized only by disturbances in motor functioning. Despite accumulating evidence for symptomatology in various non-motor domains, to date no study has investigated social cognition in CD. The aim of this study was to compare performance of CD patients and healthy controls in neurocognitive and socio-cognitive domain.

### *Methods*

Twenty-five non-depressed patients with CD and 26 healthy controls underwent neuropsychological testing. This involved assessment of cognitive status (general intellect, verbal memory, and executive function), and socio-cognitive functions using a theory of mind task and self-report on empathy and emotion regulation.

### *Results*

In comparison to controls, CD patients displayed significantly decreased cognitive abilities, particularly in executive function and verbal memory tasks. Difficulties in inferring mental states on both cognitive and affective levels were also observed. The largest discrepancies were detected in understanding intentionality in others. Poorer performance in cognitive and socio-cognitive tasks was unrelated to severity of the disease.

### *Conclusions*

This is the first evidence of compromised socio-cognitive functions in CD patients, highlighting this domain as another facet of non-motor symptoms of this disease. Future studies should advance our understanding of the extent, nature, and time course of these deficits in other aspects of social cognition in this patient population.

1    **Introduction**

2    Adult-onset idiopathic isolated cervical dystonia (CD) is a condition that is defined  
3    traditionally only by observable motor symptoms – that is, recurrent or continuous  
4    involuntary muscle contractions and subsequent abnormal postures of head and neck  
5    (Albanese et al., 2013). This relatively narrow description has been challenged over recent  
6    years, however, as evidence accumulates for a range of non-motor symptoms accompanying  
7    this disease. The most frequent findings include psychiatric comorbidities (Smit et al., 2016;  
8    Stamelou et al., 2012), alterations in sensory processing (including disease-related pain) and  
9    sensorimotor integration (Avanzino et al., 2015), sleep disturbances, and fatigue (Conte et al.,  
10   2016; Klingelhoefer et al., 2014). Importantly, some of these disturbances (e.g. depression) do  
11   not appear to be related to severity of motor difficulties (Stamelou et al., 2012) and can  
12   manifest even before onset of the disease suggesting they are an integral part of it (Conte et  
13   al., 2016). The topic is highly relevant since non-motor symptomatology (not motor  
14   difficulties as such) has a major negative impact on quality of life of dystonia patients (Smit et  
15   al., 2016; Zettenberg et al., 2012).

16       In light of the non-motor symptomatology of CD, there has been an increased effort to  
17   investigate neural abnormalities in brain areas beyond basal ganglia circuits. Existing data  
18   suggest that CD is a network disorder with several regions affected, including cerebellum,  
19   thalamus, frontal cortices, and brainstem (Battistella et al., 2015; Lehéricy et al., 2013; Zoons  
20   et al., 2011). Studies report structural abnormalities in these regions, together with alterations  
21   within various networks including abnormal hemispheric asymmetry and aberrant plasticity  
22   (Battistella et al., 2015; Quararone and Hallett, 2013). Such extensive brain changes raise the  
23   possibility of additional higher-order (or complex) cognitive capacities that could be affected  
24   in focal dystonia, and in CD specifically. Evidence for impairments in cognitive functioning  
25   is not yet conclusive (Lange et al., 2016; Stamelou et al., 2012), but subtle deficits in specific

1 domains such as attention, processing speed, and verbal memory have been reported (Kuyper  
2 et al., 2011; Romano et al., 2014; Scott et al., 2003; but see Pillon et al., 2006). This is due  
3 partly to relatively modest number of conducted studies, and also because existing data have  
4 been collected predominantly from samples that comprise different dystonia subgroups.  
5 Different forms of dystonia appear to have different phenotypes (Battistella et al., 2015;  
6 Romano et al., 2014; Scott et al., 2003), and this should be reflected in the design of studies.

7 A critical determinant for optimal functioning in our social world is having intact  
8 socio-cognitive abilities. This is true especially for a construct that is central to social  
9 cognition – Theory of Mind (ToM). ToM is defined as an ability to understand and interpret  
10 the intentions, emotions, and beliefs of others in order to predict their behaviour accurately  
11 and act accordingly (Frith and Frith, 2003). Currently, there seems to be general agreement  
12 that ToM is a multifaceted construct, encompassing cognitive (understanding intentions and  
13 beliefs) and affective components (understanding feelings and emotions; e.g. Abu-Akel &  
14 Shamay-Tsoory, 2011). The neural network subserving ToM involves posterior cortices  
15 primarily, including temporo-parietal junction, superior temporal sulcus, posterior cingulate,  
16 and precuneus. The two components of ToM can be dissociated reliably with respect to  
17 anterior and subcortical structures, however; whilst the brain circuit engaged during cognitive  
18 ToM includes dorsal parts of the anterior cingulate, medial and lateral prefrontal cortex, and  
19 the striatum, affective ToM is associated with activity in ventral aspects of these brain regions  
20 together with orbitofrontal and inferolateral frontal cortices (Abu-Akel & Shamay-Tsoory,  
21 2011; Poletti et al., 2012). Recently, there has also been an increased interest in the role of the  
22 cerebellum in ToM processing, and evidence is accumulating for its involvement in abstract  
23 mentalizing tasks (Van Overwalle & Mariën, 2016).

24 Any disturbance within this diffuse network impacts negatively on ToM abilities.  
25 Indeed, deficits in ToM have been reported in other movement disorders, including

1 Parkinson's (PD; Santangelo et al., 2012) and Huntington's disease (HD; Brüne et al., 2011;  
2 Eddy, Sira Mahalingappa, and Rickards, 2012), and essential tremor (ET; Santangelo et al.,  
3 2013). This has not been investigated in patients with idiopathic CD, however. Abnormalities  
4 found within networks involving basal ganglia-thalamo-cortical as well as cerebello-thalamo-  
5 cortical pathways in idiopathic dystonia (e.g., Lehéricy et al., 2013) should impact negatively  
6 on ToM abilities of these individuals.

7 The aim of this study was to compare performance of patients with CD and their  
8 healthy counterparts on socio-cognitive tasks, including examination of relationships with  
9 cognitive functioning and severity of the disease. Given the aforementioned neuroanatomical  
10 correlates of ToM and previous research in related populations we hypothesised that  
11 disturbances would be observed in CD patients on complex tasks and those assessing social  
12 cognition, especially its cognitive component.

13

14

1   **Methods**

2   Participants

3   The sample consisted of 25 patients with idiopathic CD (8males;  $M_{age}=56.5$ ;  $SD=12.4$ ;  
4    $M_{education}=13.3$ ;  $SD=2.8$ ) and 26 age- sex- and education-matched healthy controls (HC;  
5   7males;  $M_{age}=56.7$ ;  $SD=11.9$ ;  $M_{education}=13.3$ ;  $SD=2.6$ ). The patients were recruited from  
6   Movement Disorders Outpatient Clinic at St. Anne's University Hospital, Brno, Czech  
7   Republic. To eliminate any confounding effects of depressive symptomatology on cognitive  
8   and socio-cognitive performance, we used the Montgomery-Åsberg Depression Rating Scale  
9   (MADRS; Montgomery and Åsberg, 1979) to recruit only participants who did not suffer  
10   from clinical depression ( $MADRS \leq 16$ ; see Ketharanathan et al., 2016;  $M_{MADRS}=4.55$ ,  
11    $SD=4.10$ ). Further exclusion criteria included other forms of dystonia and additional  
12   neurological/psychiatric condition. All patients were receiving regular treatment with  
13   Botulinum toxin A for at least one year. None of the patients received further medication  
14   (anticholinergics, neuroleptics, or benzodiazepines) or deep brain stimulation treatment.  
15   Severity of the disease was assessed by experienced neurologist using Toronto Western  
16   Spasmodic Torticollis rating scale (TWSTRS; Consky et al., 1990). Average duration of CD  
17   was 12.6 years ( $SD=6.8$ ; range 1-25 years). HCs were recruited via flyers at local community  
18   centres. Exclusion criteria constituted neurological and/or psychiatric diagnosis. The study  
19   was approved by the Institutional Review Board of the St. Anne's University Hospital, and all  
20   participants provided informed consent before taking part in the study.

21

22   Materials

23   The entire neuropsychological battery was completed during the morning, in a single session  
24   that lasted approximately 3 hours.

25

1    *Cognitive abilities*  
2    Current cognitive status was assessed by selected subtests from Wechsler Adult Intelligence  
3    Scale-Revised (WAIS-R; Wechsler, 1981): Picture Completion (visual perception and  
4    attention to detail), Similarities (abstract verbal reasoning and concept formation), Digit-  
5    Symbol Coding (cognitive processing speed), and Arithmetic (working memory and  
6    numerical reasoning). The word list from Wechsler Memory Scale-III (WMS-III; Wechsler,  
7    1997) was used as a measure of verbal memory, and the interference score from Stroop test  
8    calculated according to Chafetz and Matthews (2004) was our index of cognitive control.  
9    Visual attention, task switching and planning abilities, executive control, and verbal abilities  
10   were tested by both versions of Trail Making Test (TMT; Lezak, 2004), Tower of London  
11   (Shallice, 1982), and lexical (phonemic) and semantic verbal fluency tasks (Spreen and  
12   Strauss, 2006), respectively.

13

14   *Social cognition*  
15   The Faux Pas Recognition Test (FPRT) was selected as a measure of cognitive and affective  
16   ToM due to its potential to differentiate among populations with neurodegenerative diseases  
17   (Bora, Walterfang, and Velakoulis, 2015a). The full version of the test was administered  
18   together with self-report methods on related constructs: empathy (Questionnaire of Cognitive  
19   and Affective Empathy; QCAE; Reniers et al., 2011) and emotion dysregulation (Difficulties  
20   in Emotion Regulation Scale; DERS; Gratz and Roemer, 2004). Both instruments were  
21   translated into Czech using a standard procedure of independent forward and backward  
22   translation to ensure conceptual equivalency of the items. Final versions of the translated  
23   questionnaires were then pre-tested on a large sample of healthy participants (n=255; 88  
24   males;  $M_{age}=22.5$ ;  $SD=3.1$ ), which revealed excellent levels of internal consistency (QCAE

1 cognitive:  $\alpha=.87$ ; QCAE affective:  $\alpha=.77$ ; DERS:  $\alpha=.91$ ). No difficulties with clarity or  
2 understanding were encountered with the patients themselves.

3

4           Cognitive and affective empathy

5 The QCAE includes 31 items divided into factors of cognitive and affective empathy: (1)  
6 Cognitive empathy is captured by Perspective Taking (seeing things from perspective of the  
7 others intuitively) and Online Simulation (effortful understanding of others' perspective by  
8 imagining oneself in the situation); (2) Affective empathy consists of Emotion contagion  
9 (mirroring of feelings of the others) and Proximal and Peripheral responsivity (affective  
10 responsivity in emotionally charged situations within close and detached contexts,  
11 respectively). All items are rated on a four-point Likert scale ranging from Strongly Agree to  
12 Strongly Disagree.

13

14           Emotion dysregulation

15 Over 36 items, DERS measures six aspects of emotion dysregulation: Non-acceptance of  
16 emotional responses, difficulties engaging in goal directed behaviour, impulse control  
17 difficulties, lack of emotional awareness, limited access to emotion regulation strategies, and  
18 lack of emotional clarity. Participants indicate frequency of their difficulties to regulate  
19 emotions on a scale ranging from 1 (almost never – 0-10% of time) to 5 (almost always – 91-  
20 100% of time).

21

22           Theory of Mind

23 The FPRT consists of 20 short stories describing everyday interpersonal situations.  
24 Participants listen to each story and are asked subsequently to determine whether anyone said  
25 something inappropriate or hurtful (i.e. social faux pas; FP). Ten stories include social FP

1 (FP+) and 10 control stories describe minor misunderstanding (FP-). ToM abilities are  
2 assessed using four scores: (1) FP detection (detection), (2) understanding inappropriateness  
3 of described behaviour (inappropriateness), (3) understanding intentions of story characters  
4 (intentions), (4) belief about unintentionality of FP (belief), and (5) empathy for story  
5 character affected by remark (empathy). The first and last aspect tap into affective ToM,  
6 while the rest of the scores represent cognitive ToM (Poletti, Enrici, and Adenzato, 2012). To  
7 control for the possible influence of cognitive functions of participants (e.g., attention,  
8 working memory), printed stories are available to consult for the entire testing time (without  
9 time limit). General comprehension is tested by two questions. Since four FPRT protocols did  
10 not contain data for all items, they were excluded from further analyses. FPRT responses were  
11 coded by two raters independently, with an excellent level of inter-rater reliability achieved  
12 ( $\kappa=.86$ ).

13

14 *Data analyses*

15 For subsequent analyses, raw scores from WAIS-R and WMS III subtests were transformed  
16 into standard (scaled) scores. The rest of the variables are expressed as raw scores. The FPRT  
17 scores were calculated according to the test manual (Stone, Baron-Cohen, and Knight, 1998)  
18 and interpreted in terms of separate cognitive and affective ToM (see above; Poletti, Enrici,  
19 and Adenzato, 2012). Since a majority of the variables violated assumptions of normality,  
20 performance of the two groups was compared by means of the Mann-Whitney test, and  
21 relationships between the studied variables were calculated using Spearman correlation.  
22 Effect size calculation is indicated by  $r$  (Cohen, 1998). Statistical analyses were performed in  
23 SPSS 23.

24

25 **Results**

1 *Cognition*

2 Compared with HCs, CD patients performed significantly poorer in majority of WAIS-R  
3 subtests: Picture Completion ( $U=186.50$ ;  $p=.008$ ;  $r=.37$ ), Arithmetic ( $U=203.00$ ;  $p=.021$ ;  
4  $r=.32$ ), and Digit-Symbol Coding ( $U=178.00$ ;  $p=.006$ ;  $r=.39$ ).

In addition, the CD group was significantly slower at both forms of TMT, displaying difficulties in visuomotor speed and attention (TMT A:  $U=179.50$ ;  $p=.006$ ;  $r=.38$ ) and attention distribution and cognitive flexibility (TMT B:  $U=193.00$ ;  $p=.013$ ;  $r=.35$ ). Patients' performance was also significantly worse in lexical and semantic verbal fluency ( $U=129.50$ ;  $p<.001$ ;  $r=.52$ ; and  $U=168.00$ ;  $p=.003$ ;  $r=.42$ ) and Tower of London tasks ( $U=170.50$ ;  $p=.003$ ;  $r=.41$ ). Immediate ( $U=191.00$ ;  $p=.011$ ;  $r=.36$ ) and delayed verbal memory ( $U=174.00$ ;  $p=.004$ ;  $r=.40$ ) were also significantly poorer in the patients; ability to recognize verbal material was, however, comparable in both groups ( $p=.772$ ). Similarly, no differences were found in cognitive control measured by interference score of Stroop test ( $p=.917$ ; for descriptive statistics see Table 1).

15

Table 1

17

18 *Social cognition*

19 There were no differences between the groups in self-reported cognitive and affective  
20 empathy (QCAE:  $p=.678$ ;  $p=.713$ ) and emotion regulation (DERS:  $p=.955$ ).

Actual performance in the FPRT revealed a different pattern of results, however: overall FPRT scores showed that patients performed significantly worse in FP+ ( $U=72.50$ ;  $p<.001$ ;  $r=.63$ ) as well as FP- stories ( $U=119.50$ ;  $p<.001$ ;  $r=.53$ ). In other words, the patients had difficulties not only in identifying and understanding FP accurately when it was present,

1 but also perceived inappropriate behaviour more frequently than HCs in stories, where no FP  
2 actually occurred.

3 For FP+ stories, the distinctions between CD patients and controls were significant in  
4 both cognitive (inappropriateness:  $U=114.50$ ;  $p<.001$ ;  $r=.51$ ; intentions:  $U=97.00$ ;  $p<.001$ ;  
5  $r=.56$ ; belief:  $U=112.50$ ;  $p<.001$ ;  $r=.51$ ) and affective domains (detection:  $U=120.00$ ;  $p=.001$   
6  $r=.50$ ; empathy:  $U=142.50$ ;  $p=.004$ ;  $r=.43$ ; see Figure 1).

7 Next, we analysed relationships between socio-cognitive measures and other relevant  
8 variables in our patient group. Neither cognitive nor affective empathy (as measured by  
9 QCAE) were related to FPRT performance. Emotion dysregulation indexed by DERS was,  
10 however, associated negatively with FP intentions ( $\rho=-.42$ ;  $p=.038$ ). In other words, better  
11 emotion regulation was related to a higher belief that FP was unintentional.

12 With respect to cognitive status, patients who achieved higher scores in cognitive  
13 aspects of ToM – namely, FP intentions and belief – performed significantly better in  
14 Arithmetic subtest ( $\rho=.61$ ,  $p=.001$ ;  $\rho=.55$ ;  $p=.005$ ). The former FP score was also related  
15 positively to semantic verbal fluency ( $\rho=.51$ ,  $p=.009$ ).

16 Finally, there was no relationship between severity of CD as defined by TWSTRS and  
17 any aspect of cognitive or socio-cognitive functioning ( $p>.09$ ). We observed, however,  
18 association of CD duration and FP inappropriateness ( $\rho=-.49$ ;  $p=.014$ ).

19

20 Figure1

21

22

1    **Discussion**

2    The aim of this study was to add to an existing literature on the non-motor symptomatology of  
3    idiopathic isolated CD by examining cognitive status and social cognition in this population.

4              The most important finding is a considerable decrease of performance in both aspects  
5    of ToM identified in non-depressed CD patients compared with HCs. These impairments  
6    were significant for all scores, but the largest differences were detected in cognitive  
7    component of FPRT – namely, understanding intentions of story characters. When asked  
8    about character's intentions ("Why do you think they said it?"), correct answer should express  
9    understanding that the main character did not know or realise something. The CD patients  
10   quite frequently provided answers that indicated deliberate behaviour (e.g. "He was rude and  
11   tactless.", "She was jealous and wanted to make her friend upset.", "He was showing off."),  
12   failing to realize that the character said something awkward only because they were not aware  
13   of an important detail in that particular situation. Such answers have been reported in patients  
14   with frontal lobe impairment (e.g. Gregory et al., 2002).

15             Interestingly, misinterpretation of social signals was observed in both FP+ and FP-  
16   stories. Apart from poorer understanding of individual aspects of social FP, our patients also  
17   demonstrated difficulties in interpreting minor social conflicts in otherwise neutral contexts.  
18   This cannot be interpreted as a direct consequence of poorer cognitive abilities, however,  
19   since the stories were available in front of all participants for the entire testing time.  
20   Corresponding errors have been observed in patients with other neurological and  
21   neuropsychiatric diagnoses such as HD (Eddy and Rickards, 2015), frontotemporal dementia  
22   (Gregory et al., 2002), and Asperger Syndrome (Zalla et al., 2009).

23             Poorer performance of the patient group in FPRT was in contrast to the data on self-  
24   reported cognitive and affective empathy and emotion dysregulation, where patients and  
25   controls scored similarly. This particular pattern of results – that is, a decline in both aspects

1 of ToM in patients but comparable levels in self-report – might be associated with reduced  
2 emotional awareness, interoception, or alexithymia (see Moriguchi et al., 2006). Since current  
3 knowledge on the structure of social cognition remains limited (Happé, Cook, & Bird, 2017),  
4 the relationship between higher and lower levels of socio-cognitive abilities need to be  
5 investigated further in future research. Alexithymia is often measured only via self-report  
6 (e.g., TAS-20), which might not yield reliable data owing to the possibility of poor emotional  
7 introspection in these patients. Furthermore, most recent research shows that there might be  
8 two dissociable aspects in alexithymia – affective anomia and affective agnosia (Lane et al.,  
9 2015). Any association between ToM and alexithymia should be therefore addressed using  
10 other measures such as the Levels of Emotional Awareness Scale (Lane et al., 1990) or the  
11 Modified Beth Israel Questionnaire (Taylor, Bagby, and Parker, 1997).

12 A decline in socio-cognitive abilities and ToM especially is well-documented in other  
13 movement disorders involving the basal ganglia (Poletti, Enrici, and Adenzato, 2012; Bora,  
14 Walterfang, and Velakoulis, 2015b), but the majority of published studies do not distinguish  
15 between affective and cognitive component of FPRT. Moreover, the manner of scoring of this  
16 test varies, so it is relatively problematic to determine which aspect – when considering FPRT  
17 in particular – is impaired more severely. Using different socio-cognitive tasks, existing  
18 evidence seems to point largely in the direction of cognitive ToM being affected in ET and  
19 PD (Santangelo et al., 2013; Poletti, Enrici, and Adenzato, 2012), although most recent  
20 studies show deficits also in affective ToM for PD patients (Adenzato and Poletti, 2013; Bora,  
21 Walterfang, and Velakoulis, 2015b). Additionally, both components seem to be impaired in  
22 HD (Brüne et al., 2011; Eddy, Sira Mahalingappa, and Rickards, 2012). Abnormal  
23 performance in ToM tasks manifests in neurological populations very early in the course of  
24 disease (Adenzato and Poletti, 2013; Eddy and Rickards, 2015) and tends to get more  
25 pronounced over time (Poletti, Enrici, and Adenzato 2012).

1       Regarding cognitive status, significantly worse performance of CD patients was  
2 shown in attention, set-shifting capacity, working memory, processing speed, planning, verbal  
3 memory and verbal fluency in particular. Similar patterns of cognitive impairments have been  
4 reported already in patients with focal dystonia and specifically CD (Lange et al., 2016;  
5 Romano et al., 2014). Furthermore, cognitive decline was not associated with severity of the  
6 disease, as corroborated previously by studies with similar study samples (Alemán, de  
7 Erausquin, and Micheli, 2009; Romano et al., 2013).

8       Although the relationship between social cognition and executive function in  
9 populations with neurodegenerative diseases is still a subject of ongoing discussion (Poletti  
10 Enrici, and Adenzato 2012; Adenzato and Poletti 2013; Eddy et al., 2013), some studies do  
11 report poorer social cognition in individuals with compromised executive function; for  
12 instance, verbal fluency appears to be associated with social cognition reliably in patients with  
13 PD, ET, and HD (Bora, Walterfang, and Velakoulis 2015b; Santangelo et al., 2013; Eddy,  
14 Sira Mahslingappa, and Rickards, 2012), and working memory demands seem to reduce  
15 errors made by PD patients on false-belief task (Eddy et al., 2013). This association was  
16 confirmed also in our sample, with patients' cognitive ToM related positively to working  
17 memory and verbal semantic fluency.

18       Observed results fit with the neuroanatomical correlates of CD, which points towards  
19 network-level disturbance encompassing multiple systems – not only cortico-basal-ganglia-  
20 thalamo-cortical- but also cortico-ponto-cerebello-thalamo-cortical loops with a recent focus  
21 on the role of cerebellum (Filip, Lungu, and Bareš, 2013; Lehéricy et al., 2013). Altered  
22 functioning of these large-scale circuits explains not only motor symptomatology, but also  
23 impairments in cognition and social cognition; it is becoming increasingly obvious that  
24 structures traditionally associated with motor processing such as basal ganglia and cerebellum  
25 are involved in numerous aspects of social cognition (Kemp et al., 2013; Van Overwalle &

1 Mariën, 2016). More specifically, decline in ToM as well as executive function in CD patients  
2 might be related to disturbances in fronto-striatal circuits implicated in CD (Kemp et al.,  
3 2013). We observed the largest differences in the cognitive component of ToM, which is  
4 thought to be processed by dorsolateral prefrontal cortex and dorsal striatum among other  
5 regions (Poletti, Enrici and Adenzato, 2012). This corresponds with neuroimaging data of  
6 idiopathic dystonia that points toward dysfunction in the putamen and caudate nucleus (e.g.,  
7 Lehéricy et al., 2013). Further, grey matter changes in CD were reported in the cingulate  
8 cortex and precuneus (Piccinin et al., 2015) – additional brain regions implicated in ToM  
9 processing.

10 This is the first study to report disturbances in social cognition in patients with CD.  
11 Although we have investigated a relatively small sample, we observed medium to large effect  
12 sizes for differences in cognitive and socio-cognitive abilities. Since our clinical sample was  
13 defined specifically to comprise patients with CD without clinically significant depressive  
14 symptomatology, however, the extent to which our results can be generalised to the  
15 population of CD patients in whom depression is of high incidence (e.g. Stamelou et al.,  
16 2012) or other dystonia subtypes is unknown. Secondly, because our sample included patients  
17 with a range of disease durations our data offer no insights into socio-cognitive difficulties in  
18 various stages of CD. However, our correlations suggest that socio-cognitive decline might be  
19 worsening with longer disease duration. Poor performance on the social-cognitive task we  
20 have revealed was associated with working memory and semantic verbal fluency, but not with  
21 severity of the disease. Another question worth investigating in the future is whether the  
22 decline in ToM we have observed impacts upon the quality of life in CD patients. Although  
23 depressive symptoms, disease-related pain and disability are among the most important  
24 predictors of quality of life in the CD population (Kuyper et al., 2011; Stamelou et al., 2011;  
25 Smit et al., 2016), difficulties in social interaction are also likely to contribute to worsened

1 quality of life. Since we did not administer any measure on health-related quality of life, no  
2 inferences can be made with regard to this relationship on the basis of our data. Current  
3 research in similar neurodegenerative diseases is inconclusive; while some studies report  
4 difficulties in cognitive ToM being associated with poorer quality of life (Santangelo et  
5 al.,2012), others report no such relationship (e.g., Santangelo et al., 2013). As mentioned  
6 above, our data do seem to suggest that the patients with CD might not be aware of their  
7 difficulties in social interaction. As such, it is worth investigating reports of social interactions  
8 and quality of life not only from the patients themselves, but also from their relatives or close  
9 friends.

10 Future studies are required to replicate our findings on larger samples consisting of  
11 patients with and without depressive symptomatology, and to investigate additional socio-  
12 cognitive aspects, their relationship with cognition, and the neural basis of these disturbances.  
13 Incorporating socio-cognitive tasks into routine neuropsychological assessment might help to  
14 understand challenges in every-day functioning of affected individuals and their families  
15 (Adenzato and Poletti 2013).

16

## References

- Abu-Akel A Shamay-Tsoory S. (2011) Neuroanatomical and neurochemical bases of theory of mind. *Neuropsychologia*, 49(11), 2971-2984.
- Adenzato M, Poletti M. (2013) Theory of Mind abilities in neurodegenerative diseases: An update and a call to introduce mentalizing tasks in standard neuropsychological assessments. *Clin Neuropsychiatry* 10(5):226.
- Albanese A, Bhatia K, Bressman SB, DeLong MR, Fahn S, Fung VS, ... & Lang AE. (2013) Phenomenology and classification of dystonia: a consensus update. *Mov Disord* 28(7): 863-873.
- Alemán GG, de Erausquin GA, Micheli F. (2009) Cognitive disturbances in primary blepharospasm. *Mov Disord* 24(14):2112-2120.
- Avanzino L, Tinazzi M, Ionta S, Fiorio M. (2015) Sensory-motor integration in focal dystonia. *Neuropsychologia* 79:288-300.
- Battistella G, Termsarasab P, Ramdhani RA, Fuertinger S, Simonyan K. (2015) Isolated Focal Dystonia as a Disorder of Large-Scale Functional Networks. *Cereb Cortex* bhv313.
- Bora E, Walterfang M, Velakoulis D. (2015a) Theory of mind in behavioural-variant frontotemporal dementia and Alzheimer's disease: a meta-analysis. *J Neurol Neurosurg Psychiatry* 86(7):714-719.
- Bora E, Walterfang M, Velakoulis D. (2015b) Theory of mind in Parkinson's disease: A metaanalysis. *Behav Brain Res* 292:515-520.
- Brüne M, Blank K, Witthaus H, Saft C. (2011) "Theory of mind" is impaired in Huntington's disease. *Mov Disord* 26(4):671-678.
- Cohen J. (1998) Statistical power analysis for the behavioral sciences (2nd ed.) Hillsdale, NJ: Erlbaum.
- Consky ES, Basinski A, Belle L, Ranaway R, Lang AE. (1990) The Toronto Western Spasmodic Torticollis Rating Scale (TWSTRS): assessment of validity and inter-rater reliability. *Neurology* 40(Suppl 1):445.
- Conte A, Berardelli I, Ferrazzano G, Pasquini M, Berardelli A, Fabbrini G. (2016) Non-motor symptoms in patients with adult-onset focal dystonia: sensory and psychiatric disturbances. *Parkinsonism Relat Disord* 22:S111-114.
- Chafetz MD, Matthews LH. (2004) A new interference score for the Stroop test. *Arch Clin Neuropsychol* 19(4):555-567.
- Eddy CM, Sira Mahalingappa S, Rickards HE. (2012) Is Huntington's disease associated with deficits in theory of mind?. *Acta Neurol Scand* 126(6):376-383.
- Eddy CM, Rickards HE. (2015) Theory of mind can be impaired prior to motor onset in Huntington's disease. *Neuropsychology* 29(5):792.
- Eddy CM, Beck SR, Mitchell IJ, Praamstra P, Pall HS. (2013) Theory of mind deficits in Parkinson's disease: a product of executive dysfunction?. *Neuropsychology* 27(1):37.
- Filip P, Lungu OV, Bareš M. (2013) Dystonia and the cerebellum: a new field of interest in movement disorders?. *Clin Neurophysiol* 124(7):1269-1276.
- Frith U, Frith CD. (2003) Development and neurophysiology of mentalizing. *Philos. Trans. R. Soc. Lond. Ser. B* 358, 459–473.
- Gratz KL, Roemer L. (2004) Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *J Psychopathol Behav Assess* 26(1):41-54.
- Gregory C, Lough S, Stone V, Erzinclioglu S, Martin L, Baron-Cohen S, Hodges JR. (2002) Theory of mind in patients with frontal variant frontotemporal dementia and Alzheimer's disease: theoretical and practical implications. *Brain* 125(4):752-764.

- Happé F, Cook JL, Bird G. (2017) The structure of social cognition: In (ter) dependence of sociocognitive processes. *Annual Review of Psychology*, 68, 243-267.
- Kemp J, Berthel MC, Dufour A, Despres O, Henry A, Namer IJ, Musacchio M, Sellal F. (2013) Caudate nucleus and social cognition: neuropsychological and SPECT evidence from a patient with focal caudate lesion. *Cortex* 49(2):559-571.
- Ketharanathan T, Hanwella R, Weerasundera R, de Silva, VA (2016) Diagnostic validity and factor analysis of Montgomery-Asberg Depression Rating Scale in Parkinson disease population. *Journal of geriatric psychiatry and neurology*, 29(3), pp.115-119.
- Klingelhofer L, Martino D, Martinez-Martin P, Sauerbier A, Rizos A, Jost W, Warner TT, Chaudhuri KR. (2014) Nonmotor symptoms and focal cervical dystonia: observations from 102 patients. *Basal Ganglia* 4(3):117-120.
- Kuyper DJ, Parra V, Aerts S, Okun MS, Kluger BM. (2011) Nonmotor manifestations of dystonia: a systematic review. *Mov Disord* 26(7):1206-1217.
- Montgomery SA, Åsberg, MA (1979) A new depression scale designed to be sensitive to change. *The British journal of psychiatry*, 134(4), 382-389.
- Moriguchi Y, Ohnishi T, Lane RD, Maeda M, Mori T, Nemoto K, Matsuda H, Komaki G. (2006) Impaired self-awareness and theory of mind: an fMRI study of mentalizing in alexithymia. *Neuroimage*, 32(3), 1472-1482.
- Lane RD, Quinlan DM, Schwartz GE, Walker PA, Zeitlin SB. (1990) The Levels of Emotional Awareness Scale: A cognitive-developmental measure of emotion. *Journal of personality assessment*, 55(1-2), 124-134.
- Lane RD, Hsu CH, Locke DE, Ritenbaugh C, Stonnington CM. (2015) Role of theory of mind in emotional awareness and alexithymia: Implications for conceptualization and measurement. *Consciousness and cognition*, 33, 398-405.
- Lange F, Seer C, Salchow C, Dengler R, Dressler D, Kopp B. (2016) Meta-analytical and electrophysiological evidence for executive dysfunction in primary dystonia. *Cortex* 82:133-146.
- Lehéricy S, Tijssen MA, Vidailhet M, Kaji R, Meunier S. (2013) The anatomical basis of dystonia: current view using neuroimaging. *Mov Disord* 28(7):944-957.
- Lezak MD. (2004) Neuropsychological assessment. Oxford University Press, USA.
- Pillon B, Ardouin C, Dujardin K, Vittini P, Pelissolo A, Cottencin OE, Vercueil L, Houeto JL, Krystkowiak P, Agid Y, Destée A. (2006) Preservation of cognitive function in dystonia treated by pallidal stimulation. *Neurology* 66(10):1556-1558.
- Piccinin CC, Piovesana LG, Santos MC, Guimarães RP, De Campos BM, Rezende TJ, Campos LS, Torres FR, Amato-Filho AC, França Jr MC, Lopes-Cendes I. (2015) Diffuse decreased gray matter in patients with idiopathic craniocervical dystonia: a voxel-based morphometry study. *Frontiers in neurology*, 5, p.283.
- Poletti M, Enrici I, Adenzato M. (2012) Cognitive and affective Theory of Mind in neurodegenerative diseases: neuropsychological, neuroanatomical and neurochemical levels. *Neurosci Biobehav Rev* 36(9):2147-2164.
- Quartarone A, Hallett M. (2013) Emerging concepts in the physiological basis of dystonia. *Mov Disord* 28(7):958-967.
- Reniers RL, Corcoran R, Drake R, Shryane NM, Völlm BA. (2011) The QCAE: A questionnaire of cognitive and affective empathy. *J Pers Assess* 93(1):84-95.
- Romano R, Bertolino A, Gigante A, Martino D, Livrea P, Defazio G. (2014) Impaired cognitive functions in adult-onset primary cranial cervical dystonia. *Parkinsonism Relat Disord* 20(2):162-165.
- Santangelo G, Vitale C, Trojano L, Errico D, Amboni M, Barbarulo AM, Grossi D, Barone P. (2012) Neuropsychological correlates of theory of mind in patients with early Parkinson's disease. *Mov Disord* 27:98-105.

- Santangelo G, Trojano L, Barone P, Errico D, Improta I, Agosti V, Grossi D, Sorrentino G, Vitale C. (2013) Cognitive and affective theory of mind in patients with essential tremor. *J Neurol* 260(2):513-520.
- Scott RB, Gregory R, Wilson J, Banks S, Turner A, Parkin S, Giladi N, Joint C, Aziz T. (2003) Executive cognitive deficits in primary dystonia. *Mov Disord* 18(5):539-550.
- Shallice, T. (1982) Specific impairments of planning. *Philos Trans R Soc Lond B Biol Sci* 298(1089):199–209.
- Smit M, Kuiper A, Han V, Jiawan VC, Douma G, van Harten B, et al. (2016) Psychiatric comorbidity is highly prevalent in idiopathic cervical dystonia and significantly influences health-related quality of life: Results of a controlled study. *Parkinsonism Relat Disord* 30:7-12.
- Spreen O, Strauss E. (2006) A compendium of neuropsychological tests: Administration, norms, and commentary. American Chemical Society.
- Stamelou M, Edwards MJ, Hallett M, Bhatia KP. (2012) The non-motor syndrome of primary dystonia: clinical and pathophysiological implications. *Brain* 135(6):1668-1681.
- Stone VE, Baron-Cohen S, Knight RT. (1998) Frontal lobe contributions to theory of mind. *J Cogn Neurosci* 10(5):640-656.
- Taylor GJ, Bagby RM, Parker JD. (1999) Disorders of affect regulation: Alexithymia in medical and psychiatric illness. Cambridge University Press.
- Van Overwalle F, Mariën P. (2016) Functional connectivity between the cerebrum and cerebellum in social cognition: a multi-study analysis. *NeuroImage*, 124, 248-255.
- Wechsler D. (1981) The Wechsler adult intelligence scale-revised. New York: The Psychological Corporation.
- Wechsler D. (1997) Wechsler memory scale (WMS-III). San Antonio, TX: Psychological Corporation.
- Zalla T, Sav AM, Stopin A, Ahade S, Leboyer M. (2009) Faux pas detection and intentional action in Asperger Syndrome. A replication on a French sample. *J Autism Dev Disord* 39(2):373-382.
- Zetterberg L, Lindmark B, Söderlund A, Åsenlöf P. (2012) Self-perceived non-motor aspects of cervical dystonia and their association with disability. *J Rehab Med* 44(11):950-954.
- Zoons E, Booij J, Nederveen AJ, Dijk JM Tijssen MAJ. (2011) Structural, functional and molecular imaging of the brain in primary focal dystonia—a review. *Neuroimage* 56(3):1011-1020.

**Figure 1. Differences in cognitive and affective components of FPRT**

*Note:* FPRT responses in patients (dark) and controls. Cognitive variables are highlighted; \* $p < .01$ ; \*\* $p < .001$

**Table 1. Descriptive statistic for cognitive measures**

*Note:* Parametric indices are reported here to allow comparisons with other studies.

SD=standard deviation